

ISSN 2518-1629 (Online),
ISSN 2224-5308 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
Өсімдіктердің биологиясы және биотехнологиясы институтының

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Института биологии и биотехнологии растений

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
of the Institute of Plant Biology and Biotechnology

БИОЛОГИЯ ЖӘНЕ МЕДИЦИНА
СЕРИЯСЫ



СЕРИЯ

БИОЛОГИЧЕСКАЯ И МЕДИЦИНСКАЯ



SERIES

OF BIOLOGICAL AND MEDICAL

2 (320)

НАУРЫЗ – СӘУІР 2017 ж.

МАРТ – АПРЕЛЬ 2017 г.

MARCH – APRIL 2017

1963 ЖЫЛДЫҢ ҚАҢТАР АЙЫНАН ШЫҒА БАСТАҒАН
ИЗДАЕТСЯ С ЯНВАРЯ 1963 ГОДА
PUBLISHED SINCE JANUARY 1963

ЖЫЛЫНА 6 РЕТ ШЫҒАДЫ
ВЫХОДИТ 6 РАЗ В ГОД
PUBLISHED 6 TIMES A YEAR

АЛМАТЫ, ҚР ҰҒА
АЛМАТЫ, НАН РК
ALMATY, NAS RK

Б а с р е д а к т о р

ҚР ҰҒА академигі, м. ғ. д., проф.

Ж. А. Арзықұлов

Абжанов Архат проф. (Бостон, АҚШ),
Абелев С.К. проф. (Мәскеу, Ресей),
Айтқожина Н.А. проф., академик (Қазақстан)
Ақшулаков С.К. проф., корр.-мүшесі (Қазақстан)
Алшынбаев М.К. проф., корр.-мүшесі (Қазақстан)
Березин В.Э., проф., корр.-мүшесі (Қазақстан)
Бисенбаев А.К. проф., корр.-мүшесі (Қазақстан)
Бишимбаева Н.К. проф., корр.-мүшесі (Қазақстан)
Ботабекова Т.К. проф., корр.-мүшесі (Қазақстан)
Ellenbogen Adrian prof. (Tel-Aviv, Israel),
Жамбакин К.Ж. проф., корр.-мүшесі (Қазақстан), бас ред. орынбасары
Ishchenko Alexander, prof. (Villejuif, France)
Қайдарова Д.Р. проф., корр.-мүшесі (Қазақстан)
Күзденбаева Р.С. проф., академик (Қазақстан)
Лось Д.А. prof. (Мәскеу, Ресей)
Lunefeld Bruno prof. (Израиль)
Миербеков Е.М. проф. (Қазақстан)
Муминов Т.А. проф., академик (Қазақстан)
Purton Saul prof. (London, UK)
Рахыпбеков Т.К. проф., корр.-мүшесі (Қазақстан)
Сапарбаев Мұрат проф. (Париж, Франция)
Сарбассов Дос проф. (Хьюстон, АҚШ)

«ҚР ҰҒА Хабарлары. Биология және медициналық сериясы».

ISSN 2518-1629 (Online),

ISSN 2224-5308 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5546-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
www.nauka-nanrk.kz / biological-medical.kz

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2017

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р
академик НАН РК, д.м.н., проф.

Ж. А. Арзыкулов

Абжанов Архат проф. (Бостон, США),
Абелев С.К. проф. (Москва, Россия),
Айтхожина Н.А. проф., академик (Казахстан)
Акшулаков С.К. проф., чл.-корр. (Казахстан)
Алчинбаев М.К. проф., чл.-корр. (Казахстан)
Березин В.Э., проф., чл.-корр. (Казахстан)
Бисенбаев А.К. проф., чл.-корр. (Казахстан)
Бишимбаева Н.К. проф., чл.-корр. (Казахстан)
Ботабекова Т.К. проф., чл.-корр. (Казахстан)
Ellenbogen Adrian prof. (Tel-Aviv, Israel),
Жамбакин К.Ж. проф., чл.-корр. (Казахстан), зам. гл. ред.
Ishchenko Alexander prof. (Villejuif, France)
Кайдарова Д.Р. проф., чл.-корр. (Казахстан)
Кузденбаева Р.С. проф., академик (Казахстан)
Лось Д.А. prof. (Москва, Россия)
Lunenfeld Bruno prof. (Израиль)
Миербеков Е.М. проф. (Казахстан)
Муминов Т.А. проф., академик (Казахстан)
Purton Saul prof. (London, UK)
Рахыпбеков Т.К. проф., чл.-корр. (Казахстан)
Сапарбаев Мурат проф. (Париж, Франция)
Сарбассов Дос проф. (Хьюстон, США)

«Известия НАН РК. Серия биологическая и медицинская».

ISSN 2518-1629 (Online),

ISSN 2224-5308 (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов
Министерства культуры и информации Республики Казахстан №5546-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел. 272-13-19, 272-13-18,
www.nauka-nanrk.kz/biological-medical.kz

© Национальная академия наук Республики Казахстан, 2017

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

Editor in chief

academician of NAS RK, doctor of medical science, professor

Zh. A. Arzykulov

Abzhanov Arkhat prof. (Boston, USA),
Abelev S.K. prof. (Moscow, Russia),
Aitkhozhina N.A. prof., academician (Kazakhstan)
Akshulakov S.K. prof., corr. member. (Kazakhstan)
Alchinbayev M.K. prof., corr. member. (Kazakhstan)
Berezin V.Ye., prof., corr. member. (Kazakhstan)
Bisenbayev A.K. prof., corr. member. (Kazakhstan)
Bishimbayeva N.K. prof., corr. member. (Kazakhstan)
Botabekova T.K. prof., corr. member. (Kazakhstan)
Ellenbogen Adrian prof. (Tel-Aviv, Israel),
Zhambakin K.Zh. prof., corr. member. (Kazakhstan), deputy editor in chief
Ishchenko Alexander, prof. (Villejuif, France)
Kaydarova D.R. prof., corr. member. (Kazakhstan)
Kuzdenbayeva R.S. prof., academician (Kazakhstan)
Los D.A. prof. (Moscow, Russia)
Lunefeld Bruno prof. (Israel)
Miyerbekov Ye.M. prof. (Kazakhstan)
Muminov T.A. prof., academician (Kazakhstan)
Purton Saul prof. (London, UK)
Rakhypbekov T.K. prof., corr. member. (Kazakhstan)
Saparbayev Murat prof. (Paris, France)
Sarbassov Dos, prof. (Houston, USA)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of biology and medicine.

ISSN 2518-1629 (Online),

ISSN 2224-5308 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5546-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,

<http://nauka-nanrk.kz/biological-medical.kz>

© National Academy of Sciences of the Republic of Kazakhstan, 2017

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF BIOLOGICAL AND MEDICAL

ISSN 2224-5308

Volume 2, Number 320 (2017), 21 – 26

UDC 546.15+57.085.23+615.28+59.085

K. Bekesheva^{1,2}, N. Zubenko², G. Kon², T. Kustova²,
R. Islamov², G. Ustenova¹, T. Baczek³, A. Ilin²

¹S. D. Asfendiyarov Kazakh National Medical University, Almaty, Kazakhstan,

²Scientific Center for Anti-Infectious Drugs, Almaty, Kazakhstan,

³Department of Pharmaceutical Chemistry, Medical University of Gdansk, Gdansk, Poland.

E-mail: renatislamov@gmail.com

CYTOTOXICITY AND ACUTE TOXICITY OF A NEW COMPOUND COMPRISING IODINE ADDUCTS IN MICE

Abstract. The cytotoxicity and acute toxicity were studied after oral and intraperitoneal administration of a new compound (R8) comprising iodine adducts to mice. The median cytotoxic concentration (CC₅₀) was determined in the six cell lines, and ranged from 2.1 to 3.9 mg/ml. The median lethal dose of the compound R8 in mice when administered orally was 1000 mg/kg, intraperitoneally – 200 mg/kg. According to the international system of classification for toxicity of chemicals GHS, the compound R8 may be assigned to Toxicity Category IV.

Keywords: iodine, adducts, cytotoxicity, acute toxicity, mice.

Introduction. The prevalence of resistance to colistin and carbapenems in bacteria actually disarms the modern anti-infectious therapy [1]. For this reason, the urgent is the search for new approaches and strategies for the treatment of drug-resistant infections [2]. Currently, iodine is among the few substances that have retained antimicrobial properties [3-6]. The preparations comprising complexes of iodine with organic macromolecules such as povidone-iodine and cadexomers, have become widespread in human and veterinary medicine [5, 7]. In addition, the complexes of iodine with dextran increase expression of IL-1 β , IL-8, TNF- α , and VEGF mRNA and do not affect IL-6, IL-10, IL-12 and bFGF mRNA, which indicates the specific immunotropic activity [8]. At the same time, their use is limited to relatively high chemical activity and toxicity of povidone-iodine [7,9]. One of the ways to reduce iodine toxicity lies in its combining with carbohydrates and producing a complex with dextrin (cadexomers). A comparative study on the effect of various complex compounds of iodine on the blood vessels in the chick embryo chorioallantoic membrane showed that povidone-iodine at 1% concentration is characterized by strong irritating activity, whereas cadexomer at a concentration up to 1.8% had a moderate activity with an exposure for 60 minutes [9]. This approach was used in the synthesis of a new stable iodine-containing substance. A new coordination iodine compound (R8) has been synthesized in the Scientific Center for Anti-Infectious Drugs [10]. To carry out initial assessment of toxicity of the compound R8, the cytotoxicity was studied in various mammalian cell cultures, and acute toxicity studies were performed in mice.

Materials and Methods. *Test substance.* Coordination iodine compound with molecular weight of 458.11 and the chemical formula C₁₈H₂₃I₃N₂O₄. The substance at 20°C and 101.3 kPa is a solid crystalline gray-green powder with a faint iodine odor. It is readily soluble in water. Immediately before performing assays, the R8 substance was dissolved in sterile water.

Cell cultures. RD - human embryonic rhabdomyosarcoma cells, BHK-21 - newborn Syrian hamster kidney fibroblasts, MDCK - dog kidney cell line, Hep G2 - human hepatocellular carcinoma cells (ATCC). The subinoculated suspension cell cultures: H9 – human cutaneous T-cell lymphoma (ATCC), K562 - human erythroblastoid leukemic cells (ATCC).

Laboratory animals. White female outbred mice in the number of 24 individuals, aged 1.5-2.5 months and weighing $22 \text{ g} \pm 10\%$, were obtained from the Scientific and Practical Center for Sanitary-Epidemiological Expertise and Monitoring (Almaty, Kazakhstan). The mice were kept in individually ventilated cages (IVC) (Tecniplast, Italy) in controlled conditions (with a 12-hour day/night cycle, at a temperature of $22 \pm 2^\circ\text{C}$ and relative humidity of $55 \pm 10\%$). The mice were fed the commercial complete feed (“Assortiment-Agro” LLC, Russia). After a 2-week acclimatization period, the animals were randomized into groups. The animals were given *ad libitum* access to food and water.

Cytotoxicity. Cytotoxicity was evaluated by analyzing the cell’s ability of absorbing neutral red in accordance with the OECD Series on Testing and Assessment, No. 129 “Guidance document on using cytotoxicity tests to estimate starting doses for acute oral systemic toxicity tests”. The cells were seeded in 96-well plates at a concentration of 2.5×10^5 cells in 1.0 ml. The plates were incubated in a 5% CO₂ incubator at 37°C. After incubation for 24 hours, the culture medium was removed from the wells, and 200.0 µl of the incomplete medium containing the test substance R8 were added to each well. The R8 concentrations were prepared by double dilution with DMEM medium (for HepG2, RD, 21-BHK, MDCK cell cultures) or with RPMI-1640 medium (for H9, R562 cell cultures), starting from 10.0 mg/ml. 200.0 µl of incomplete culture medium (RPMI-1640 for H9, K562 and DMEM for HepG2, RD, 21-BHK, MDCK) were added to each negative control well. For HepG2, RD, BHK-21 or MDCK cell cultures, the culture medium with the substance was removed from the wells after 48 and 72 hours, and 150.0 µl of Dulbecco’s Phosphate Buffered Saline (DPBS) were added to each well (to wash the cells). After completion of the washing step, the supernatant was removed from the plate, and 100.0 µl of the neutral red working solution were added to each well. The plate was incubated for 3 hours at 37°C. After incubation of the plate, the supernatant was removed and 150.0 µl of DPBS was added (to wash the cells). Further, the supernatant was removed from the plate, and 150.0 µl of lysis solution (mixture of 1% glacial acetic acid, 50% ethanol and 49% distilled water) were added to each well. For K562 or H9 cell cultures, the plate was centrifuged after 48 and 72 hours at 130 g for 5 minutes; the culture medium with the substance was further removed from the wells of the plate, and 250.0 µl of the neutral red working solution were added to each well. The plate was incubated for 3 hours at 37°C. After incubation, the plate was centrifuged at 130 g for 5 minutes; the supernatant was further removed from the wells of the plate, and 250.0 µl of the washing/fixing solution (0.5% formaldehyde, 99.5% distilled water) were added to each well. The plate was centrifuged at 130 g for 2 min, the supernatant was removed, and 100.0 µl of lysis solution were added. The optical density in the wells was measured with a microplate reader Tecan Sunrise RC.4 (Austria) at a wavelength of the main filter of 540 nm. Evaluation of the results and calculation of the median cytotoxic concentration (CC₅₀) were carried out based on the methods described in [11, 12].

Acute toxicity. Acute toxicity was evaluated according to the OECD Guidelines for the Testing of Chemicals, Test No. 423: Acute oral toxicity - Acute toxic class method. The test substance R8 was administered intragastrically and intraperitoneally at doses of 50, 300, and 2000 mg/kg. Toxicity category was assigned for the compound R8 as indicated in the international classification system for toxicity of chemicals GHS, according to Table 1.

The body weight of animals was measured weekly, starting on the first day of the study and until euthanasia following the completion of the experiment. The clinical signs of poisoning were recorded. The mice were killed by cervical dislocation. At necropsy, a macroscopic examination of organs was performed. The experiments on animals were approved by the Ethics Committee of the Scientific Center (No. 04-03 037) dated 26.1.2016, and Local Ethics Committee of the S.D. Asfendiyarov KazNMU dated 24.02.2016 (Protocol No. 2).

Table 1 – GHS classification of toxic chemicals

Toxicity category	Category 1	Category 2	Category 3	Category 4	Category 5
LD ₅₀ , mg/kg	< 5	> 5 < 50	> 50 < 300	> 300 < 2000	> 2000 < 5000

Statistical data processing. The means and standard deviations were calculated for the body weight of mice. The Mann-Whitney test for significance level at $p < 0.05$ was used to test the null hypothesis.

Results and Discussion. *Cytotoxicity.* The results of cytotoxicity evaluations for R8 on various cell cultures after 48 and 72 hours of incubation are shown in Table 2.

Table 2 – Results of determining cytotoxic effects of R8 in the neutral red test

Cell culture	CC ₅₀ , mg/ml	
	48 h	72 h
MDCK	3.8	3.9
HepG2	3.7	3.6
RD	2.4	2.1
BHK-21	2.1	2.1
H9	3.1	2.3
K562	2.2	1.05

Despite rather wide range of variability in the results of determining CC₅₀ for R8 within 2.1 to 3.8 mg/ml after 48 hours of incubation, these concentrations are close to 5 mg/mL (or 10 mM), which indicates the relative nontoxicity of the compound [13]. For comparison, the concentration of povidone-iodine at which 50% of murine fibroblasts (L929) survive after a 30-min incubation in the neutral red assay is 460 - 490 mg/ml or 1.8 - 1.9 mM of iodine [14]. Thereby, the new iodine adduct exhibits lower cytotoxicity as compared with povidone-iodine, even with long-lasting incubation. To evaluate the R8 safety, the acute toxicity was examined using two routes of administration, enteral and parenteral.

Acute toxicity after the intragastric administration. After administration of an initial dose of R8 (300 mg/kg), none of the animals died within 14 days. In the absence of a lethal effect, the administered dose was increased to 2000 mg/kg. After the intragastric administration of the R8 substance at a dose of 2000 mg/kg, one mouse died after 24 hours and one more mouse died 48 hours later. In total, 2 mice died in the experiment. The data obtained are shown in Table 3.

Table 3 – Mortality of mice after the intragastric administration of R8

Groups	Dose, mg/kg	Effect (died/initial number)	Time of death
Control	–	0/3	–
Experimental 1	300	0/6	–
Experimental 2	2000	2/3	Days 1 and 2

To evaluate the adverse effect of R8 in mice from the group with the dose of 300 mg/kg, the dynamics of body weight was studied in the experiment (Table 4).

Table 4 – Dynamics of body weight in mice after the intragastric administration of R8

Groups	Dose	Average weight of animals, g	
		Day 1	Day 14
Control	Solvent (water)	19.80±0.35	24.7±1.8
Experimental 1	300 mg/kg	21.4±0.75	25.2±1.1

The positive dynamics of body weight and absence of clinical signs to confirm toxic effect after exposure to 300 mg/kg suggest that LD₅₀ for R8 is about 1000 mg/kg.

On day 14 the animals were killed by cervical dislocation and subject to necropsy. The macroscopic examination of internal organs of experimental animals did not reveal any pathological deviations. At the same time, the arrangement of internal organs was normal; no cohesion between them or sharp increase or decrease in size were observed. The liver was deep red, with smooth capsule, homogeneous on the cut surface, the consistency was normal. Hemorrhages or fat deposits in the liver parenchyma were not detected. The loops of the small intestine were free. The walls of the intestine and bladder were not damaged.

The renal capsule was easily removed. The spleen was deep cherry in color, does not produce scrape-off on the cut surface, non-ulcerated and without signs of hemorrhages. The heart muscles were dark red, homogeneous, free of injuries. Hemorrhages were not detected. The lung tissue was airy, pink, with no signs of edema, hemorrhages or necrosis on the cut surface.

Acute toxicity after the intraperitoneal administration. A single-dose intraperitoneal administration of R8 at a dose of 300 mg/kg on day 3 resulted in the death of one mouse (Table 5). Immediately after the administration of R8, the animals bunched up. Within two days before the death, the mice developed arched back and exhibited a significant reduction in motor activity. Thereafter, the dose of R8 was reduced to 50 mg/kg. After administration of R8 at a dose of 50 mg/kg the mice bunched up within two hours. These symptoms gradually disappeared on the third hour. Mortality was not observed (Table 5).

Table 5 – Mortality of mice after the intraperitoneal administration of R8

Groups	Dose, mg/kg	Effect (died/initial number)	Time of death
Control	–	0/3	–
Experimental 1	300	1/3	Day 3
Experimental 2	50	0/6	–

The positive dynamics of body weight (Table 6) and the observed clinical signs of toxicity allow us to ascertain that the LD₅₀ for R8 after intraperitoneal administration is about 200 mg/kg.

Table 6 – Dynamics of body weight in mice after the intraperitoneal administration of R8

Groups	Dose	Average weight of animals, g	
		Day 1	Day 14
Control	Solvent (water)	23.0±0.4	27.2±1.0
Experimental 1	50 mg/kg	23.5±0.4	25.4±1.6

All surviving mice were killed by cervical dislocation on day 14 and subject to necropsy. The macroscopic examination of internal organs of experimental animals did not reveal any pathological deviations. When examining site for injecting R8 in the right lower third of the stomach, the inflammation and commissural processes were not found. The arrangement of internal organs was normal; no cohesion between them or sharp increase or decrease in size were observed.

Correlation between the cytotoxicity and acute toxicity data. Currently, various methodological approaches were developed for using cytotoxicity test in predicting the experiments on laboratory animals. The results can be applied to the oral acute toxicity studies using oral administration, for example, to calculate the initial dose [15]. Using the regression equation for IC₅₀ – LD₅₀ [15] from the data on cytotoxicity for R8 (2.1 mg/ml), the expected dose was evaluated in an *in vivo* experiment. As a result, the estimated LD₅₀ for R8 is 1819 mg/kg, which is higher than the experimental dose of 1000 mg/kg. These differences may be due to the mechanisms of toxic action of the compound R8.

Conclusion. The median lethal dose of the new iodine adduct R8 in mice when administered orally is 1000 mg/kg, intraperitoneally - 200 mg/kg. At that, the examined iodine adduct was less toxic as compared with iodophor - povidone-iodine. According to the international classification system for toxicity of chemicals GHS, the compound R8 may be assigned to Toxicity Category IV.

REFERENCES

- [1] Qureshi ZA, Hittle LE, O'Hara JA, Rivera JI, Syed A, Shields RK, Pasculle AW, Ernst RK, Doi Y (2015) Colistin-resistant *Acinetobacter baumannii*: beyond carbapenem resistance, Clin Infect Dis, 60:1295–1303. DOI: 10.1093/cid/civ048.
- [2] Marston HD, Dixon DM, Knisely JM, Palmore TN, Fauci AS (2016) Antimicrobial resistance, JAMA, 316:1193-1204. DOI: 10.1001/jama.2016.11764
- [3] Cooper RA (2007) Iodine revisited, Int Wound J, 4. P:124-137. DOI: 10.1111/j.1742-481X.2007.00314.x
- [4] Zubko EI, Zubko MK (2013) Co-operative inhibitory effects of hydrogen peroxide and iodine against bacterial and yeast species, BMC Res Notes, 15:272. DOI: 10.1186/1756-0500-6-272

- [5] Kanagalingam J, Feliciano R, Hah JH, Labib H, Le TA, Lin JC (2015) Practical use of povidone-iodine antiseptic in the maintenance of oral health and in the prevention and treatment of common oropharyngeal infections, *Int J Clin Pract*, 69:1247-56. DOI: 10.1111/ijcp.12707
- [6] Gottardi W, Debatov D, Nagl M (2013) N-chloramines, a promising class of well-tolerated topical anti-infectives, *Antimicrob Agents Chemother*, 57:1107-1114. DOI: 10.1128/AAC.02132-12
- [7] Noda Y, Fujii K, Fujii S (2009) Critical evaluation of cadexomer-iodine ointment and povidone-iodine sugar ointment, *Int J Pharm*, 372:85-90. DOI: 10.1016/j.ijpharm.2009.01.007
- [8] Ohtani T, Mizuashi M, Ito Y, Aiba S (2007) Cadexomer as well as cadexomer iodine induces the production of proinflammatory cytokines and vascular endothelial growth factor by human macrophages, *Exp Dermatol*, 16:318-323. DOI: 10.1111/j.1600-0625.2006.00532.x
- [9] Marquardt C, Matuschek E, Bölke E, Gerber PA, Peiper M, Seydlitz-Kurzbach J, Bühren BA, van Griensven M, Budach W, Hassan M, Kukova G, Mota R, Höfer D, Orth K, Fleischmann W (2010) Evaluation of the tissue toxicity of antiseptics by the hen's egg test on the chorioallantoic membrane (HETCAM), *Eur J Med Res*, 15:204-209. DOI: 10.1186/2047-783X-15-5-204.
- [10] Karzhaubayeva R, Bekesheva K, Kalykova A, Kabdrasova A, Ustenova G, Barinov D (2016) Standardization method of quantitative determination of iodide ions in substances and tablets containing iodine adducts, *RJPBCS*, 7:1639-1644
- [11] Repetto G, del Peso A, Zurita J L (2008) Neutral red uptake assay for the estimation of cell viability/cytotoxicity, *Nature Protocols*, 3:1125-1131. DOI: 10.1038/nprot.2008.75
- [12] Dragonov M (2004) Kletochni kul'tury. Teorija i tehniki. VAP, Plovdiv. (In Bulgarian)
- [13] Gad SC (2000) *In vitro toxicology*, second edition. Taylor and Francis, N.-Y. ISBN:1560327693
- [14] Müller G, Kramer A (2006) Comparative study of in vitro cytotoxicity of povidone-iodine in solution, in ointment or in a liposomal formulation (Repithel) and selected antiseptics, *Dermatology*, 212:91-93. DOI: 10.1159/000090102
- [15] OECD Environment, Health and Safety Publications Series on Testing and Assessment No. 129. Guidance document on using cytotoxicity tests to estimate starting doses for acute oral systemic toxicity tests. Paris, OECD, 2010.

Қ. Бекешева^{1,2}, Н. Зубенко², Г. Кон², Т. Кустова², Р. Исламов^{2*}, Г. Устенова¹, Т. Вазек³, А. Ильин²

¹Асфендияров атындағы Қазақ ұлттық медициналық университеті, Алматы, Қазақстан,

²Инфекцияға қарсы препараттар ғылыми орталығы, Алматы, Қазақстан,

³Фармацевтикалық химия факультеті, Гданьск медициналық университеті, Гданьск, Польша

ҚҰРАМЫНДА ИОД АДДУКТЫ БАР ЖАҢА ҚОСЫЛЫСТЫҢ ТЫШҚАНДАРҒА ЦИТОУЫТТЫЛЫҒЫ МЕН ӨТКІР УЫТТЫЛЫҒЫ

Аннотация. Құрамында иод аддуктысы бар жаңа қосылыстың (R8) тышқандарға ауызынан және перитонеальдық тәсілмен енгізгенде цитоуттылығы мен өткір уыттылығы зерттелінген. Орташа цитоуыттылық концентрациясы жасушалардың алты түрінде анықталды және 2,1 ден 3,9 мг/мл ауытқу аралығында болды. Тышқандардың орташа өлім дозасы R8 ауызынан енгізгенде 1000 мг/кг, ал перитонеальдық тәсілмен енгізгенде 200 мг/кг құрады. GHS халықаралық дәрілік заттардың уыттылық классификациясы жүйесіне сәйкес R8 қосылысын уыттылықтың 4 класына жатқызуға болады.

Түйін сөздер: иод, аддукт, цитоуыттылық, өткір уыттылық, тышқандар.

К. Бекешева^{1,2}, Н. Зубенко², Г. Кон², Т. Кустова², Р. Исламов², Г. Устенова¹, Т. Вазек³, А. Ильин²

¹Казахский Национальный медицинский университет им. Асфендиярова, Алматы, Казахстан,

²Научный центр противоионфекционных препаратов, Алматы, Казахстан,

³Department of pharmaceutical chemistry, Medical University of Gdansk, Gdansk, Poland

ЦИТОТОКСИЧНОСТЬ И ОСТРАЯ ТОКСИЧНОСТЬ НА МЫШАХ НОВОГО СОЕДИНЕНИЯ СОДЕРЖАЩЕГО АДДУКТЫ ИОДА

Аннотация. Изучена цитотоксичность и острая токсичность при оральном и внутрибрюшинном способах введения мышам нового соединения (R8) содержащего аддукты иода. Средняя цитотоксическая концентрация (CC₅₀) определена на шести линиях клеток и находилась в пределах от 2,1 до 3,9 мг/мл. Средняя смертельная доза на мышах при оральном поступлении R8 составила 1000 мг/кг, при внутрибрюшинном 200 мг/кг. Согласно международной системе классификации токсичности веществ GHS соединение R8 можно отнести 4 классу токсичности.

Ключевые слова: иод, аддукты, цитотоксичность, острая токсичность, мышы.

Сведения об авторах:

Бекешева К.Б. – PhD-докторант КазНМУ им. С. Д. Асфендиярова, управляющий отделом обеспечения качества GMP АО «Научный центр противоиных препаратов», e-mail: arun82@bk.ru

Зубенко Н. – старший научный сотрудник лаборатории вирусологии АО «Научный центр противоиных препаратов» e-mail: z.lyudmila_@mail.ru

Кон Г.А. – научный сотрудник лаборатории фармакологии и токсикологии, АО «Научный центр противоиных препаратов»

Кустова Т.С. – заведующий лабораторией фармакологии и токсикологии, АО «Научный центр противоиных препаратов», PhD

Исламов Р.А. – начальник отдела доклинических испытаний, АО «Научный центр противоиных препаратов», кандидат биологических наук, e-mail: renatislamov@gmail.com

Устенова Г.О. – директор учебного департамента фармации Казахского национального медицинского университета им. С. Д. Асфендиярова, доктор фармацевтических наук, доцент e-mail: ustenova@list.ru

Baczek T. – department of Pharmaceutical Chemistry, Medical University of Gdansk, Poland. e-mail: tbaczek@gumed.edu.pl

Ильин А.И. – председатель Правления АО «Научный центр противоиных препаратов», доктор химических наук, академик КазНАЕН.

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www.nauka-nanrk.kz

ISSN 2518-1629 (Online), ISSN 2224-5308 (Print)

<http://www.biological-medical.kz/index.php/ru/>

Редактор *М. С. Ахметова, Д. С. Аленов, Т. М. Апендиев*
Верстка на компьютере *Д. Н. Калкабековой*

Подписано в печать 06.04.2017.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
16,4 п.л. Тираж 300. Заказ 2.