

ISSN 2518-1629 (Online),
ISSN 2224-5308 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
Өсімдіктердің биологиясы және биотехнологиясы институтының

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Института биологии и биотехнологии растений

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
of the Institute of Plant Biology and Biotechnology

**БИОЛОГИЯ ЖӘНЕ МЕДИЦИНА
СЕРИЯСЫ**



СЕРИЯ

БИОЛОГИЧЕСКАЯ И МЕДИЦИНСКАЯ



SERIES

OF BIOLOGICAL AND MEDICAL

3 (321)

МАМЫР – МАУСЫМ 2017 ж.

МАЙ – ИЮНЬ 2017 г.

MAY – JUNE 2017

1963 ЖЫЛДЫҢ ҚАҢТАР АЙЫНАН ШЫҒА БАСТАҒАН
ИЗДАЕТСЯ С ЯНВАРЯ 1963 ГОДА
PUBLISHED SINCE JANUARY 1963

ЖЫЛЫНА 6 РЕТ ШЫҒАДЫ
ВЫХОДИТ 6 РАЗ В ГОД
PUBLISHED 6 TIMES A YEAR

АЛМАТЫ, ҚР ҰҒА
АЛМАТЫ, НАН РК
ALMATY, NAS RK

Б а с р е д а к т о р

ҚР ҰҒА академигі, м. ғ. д., проф.

Ж. А. Арзықұлов

Абжанов Архат проф. (Бостон, АҚШ),
Абелев С.К. проф. (Мәскеу, Ресей),
Айтқожина Н.А. проф., академик (Қазақстан)
Акшулаков С.К. проф., корр.-мүшесі (Қазақстан)
Алшынбаев М.К. проф., корр.-мүшесі (Қазақстан)
Березин В.Э., проф., корр.-мүшесі (Қазақстан)
Бисенбаев А.К. проф., корр.-мүшесі (Қазақстан)
Бишимбаева Н.К. проф., корр.-мүшесі (Қазақстан)
Ботабекова Т.К. проф., корр.-мүшесі (Қазақстан)
Ellenbogen Adrian prof. (Tel-Aviv, Israel),
Жамбакин К.Ж. проф., корр.-мүшесі (Қазақстан), бас ред. орынбасары
Ishchenko Alexander, prof. (Villejuif, France)
Қайдарова Д.Р. проф., корр.-мүшесі (Қазақстан)
Күзденбаева Р.С. проф., академик (Қазақстан)
Лось Д.А. prof. (Мәскеу, Ресей)
Lunefeld Bruno prof. (Израиль)
Миербеков Е.М. проф. (Қазақстан)
Муминов Т.А. проф., академик (Қазақстан)
Purton Saul prof. (London, UK)
Рахыпбеков Т.К. проф., корр.-мүшесі (Қазақстан)
Сапарбаев Мұрат проф. (Париж, Франция)
Сарбассов Дос проф. (Хьюстон, АҚШ)

«ҚР ҰҒА Хабарлары. Биология және медициналық сериясы».

ISSN 2518-1629 (Online),

ISSN 2224-5308 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5546-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
www.nauka-nanrk.kz / biological-medical.kz

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2017

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р
академик НАН РК, д.м.н., проф.

Ж. А. Арзыкулов

Абжанов Архат проф. (Бостон, США),
Абелев С.К. проф. (Москва, Россия),
Айтхожина Н.А. проф., академик (Казахстан)
Акшулаков С.К. проф., чл.-корр. (Казахстан)
Алчинбаев М.К. проф., чл.-корр. (Казахстан)
Березин В.Э., проф., чл.-корр. (Казахстан)
Бисенбаев А.К. проф., чл.-корр. (Казахстан)
Бишимбаева Н.К. проф., чл.-корр. (Казахстан)
Ботабекова Т.К. проф., чл.-корр. (Казахстан)
Ellenbogen Adrian prof. (Tel-Aviv, Israel),
Жамбакин К.Ж. проф., чл.-корр. (Казахстан), зам. гл. ред.
Ishchenko Alexander prof. (Villejuif, France)
Кайдарова Д.Р. проф., чл.-корр. (Казахстан)
Кузденбаева Р.С. проф., академик (Казахстан)
Лось Д.А. prof. (Москва, Россия)
Lunenfeld Bruno prof. (Израиль)
Миербеков Е.М. проф. (Казахстан)
Муминов Т.А. проф., академик (Казахстан)
Purton Saul prof. (London, UK)
Рахыпбеков Т.К. проф., чл.-корр. (Казахстан)
Сапарбаев Мурат проф. (Париж, Франция)
Сарбассов Дос проф. (Хьюстон, США)

«Известия НАН РК. Серия биологическая и медицинская».

ISSN 2518-1629 (Online),

ISSN 2224-5308 (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов
Министерства культуры и информации Республики Казахстан №5546-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел. 272-13-19, 272-13-18,

www.nauka-nanrk.kz/biological-medical.kz

© Национальная академия наук Республики Казахстан, 2017

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

Editor in chief

academician of NAS RK, doctor of medical science, professor

Zh. A. Arzykulov

Abzhanov Arkhat prof. (Boston, USA),
Abelev S.K. prof. (Moscow, Russia),
Aitkhozhina N.A. prof., academician (Kazakhstan)
Akshulakov S.K. prof., corr. member. (Kazakhstan)
Alchinbayev M.K. prof., corr. member. (Kazakhstan)
Berezin V.Ye., prof., corr. member. (Kazakhstan)
Bisenbayev A.K. prof., corr. member. (Kazakhstan)
Bishimbayeva N.K. prof., corr. member. (Kazakhstan)
Botabekova T.K. prof., corr. member. (Kazakhstan)
Ellenbogen Adrian prof. (Tel-Aviv, Israel),
Zhambakin K.Zh. prof., corr. member. (Kazakhstan), deputy editor in chief
Ishchenko Alexander, prof. (Villejuif, France)
Kaydarova D.R. prof., corr. member. (Kazakhstan)
Kuzdenbayeva R.S. prof., academician (Kazakhstan)
Los D.A. prof. (Moscow, Russia)
Lunefeld Bruno prof. (Israel)
Miyerbekov Ye.M. prof. (Kazakhstan)
Muminov T.A. prof., academician (Kazakhstan)
Purton Saul prof. (London, UK)
Rakhypbekov T.K. prof., corr. member. (Kazakhstan)
Saparbayev Murat prof. (Paris, France)
Sarbassov Dos, prof. (Houston, USA)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of biology and medicine.

ISSN 2518-1629 (Online),

ISSN 2224-5308 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5546-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/biological-medical.kz>

© National Academy of Sciences of the Republic of Kazakhstan, 2017

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF BIOLOGICAL AND MEDICAL

ISSN 2224-5308

Volume 3, Number 321 (2017), 39 – 43

UDC 628.336.6

K. U. Sultangaliyeva, Zh. K. Bakhov, A. A. Abubakirova, B. Zh. Mutaliyeva

M. Auezov South Kazakhstan State University, Shymkent, Kazakhstan.

E-mail: swallow0101@mail.ru, zhbakhov@mail.ru, azhar.baikal79@mail.ru, m_bota@mail.ru

**EVALUATION OF THE ANAEROBIC DIGESTION
OF AGRICULTURAL WASTE IN LEACH-BED REACTOR
FOR BIOGAS PRODUCTION**

Abstract. Anaerobic digestion processes for agricultural wastes are widely used to reduce environmental and social issues and as a way of alternative renewable energy production. Anaerobic dry fermentation of agricultural waste (cattle manure) was investigated to evaluate an efficiency of the single-stage leach-bed process for biogas production. The experiments were performed in batch-operation mode at the temperature of $40 \pm 0,2^\circ\text{C}$. Bioreactor with a working volume of 50 L had been constructed by modifying leach-bed reactor and fixed-bed reactor. The reactor was equipped with immobilization device positioned at the bottom as a layer of polyethylene packing rings. Experimental reactors were tested during a period of 28 days in Run 1 and 21 days in Run 2. The performance of the reactor was analyzed in terms of the biogas production. According to the results, the average cumulative biogas yield was $(0.331 \pm 0.005) \text{ Nm}^3 (\text{kg oDM})^{-1}$, average percentage of methane was $(47.13 \pm 1.40)\%$ during the anaerobic digestion. Anaerobic digestion of agricultural waste in leach-bed reactor is feasible and stable process for biogas production without anaerobic pretreatment and mixing, and immobilization of microflora on the supporting material improved methane production.

Keywords: anaerobic digestion, biogas, agricultural waste, methane, leach-bed reactor.

Introduction. Nowadays, anaerobic digestion processes for agricultural wastes are widely used to reduce environmental and social issues in the areas of livestock and poultry farms, and it has become an alternative way of renewable energy production. Numerous studies had been conducted in order to increase and stabilize biogas yield, optimize anaerobic biogas technologies. One of the ways of optimization of biogas production is improvement of bioreactors design. Since successful application of anaerobic fermentation technologies for treatment of organic waste from agricultural and industrial production is largely dependent on the development and use of high-rate anaerobic bioreactors. In such reactors, a large amount of substrate is processed, and optimally designed bioreactor can reduce processing time and improve the processing efficiency, resulting in an overall reduction in processing cost [1].

And such high-rate reactors as the anaerobic stirred tank reactor, the anaerobic contact process, the anaerobic sequentially cyclic reactor, the anaerobic fixed-bed reactor, the expanded or fluidized-bed reactor, the upflow anaerobic sludge blanket reactor, the anaerobic barrier reactor, leach-bed reactor can be mentioned [2;3]. In such reactors, the average growth rate of methanogenic bacteria is much lower than the growth rate of acidogenic bacteria and the overall rate of biomethanogenesis process is controlled by methanogenesis stage in anaerobic fermentation of waste water with the low solids content. Parawira (2004) [4] in his works found out, that the biomethanogenesis rate can be accelerated by increase of conversion rate of VFA into methane by increasing the concentration of the methanogenic bacteria in the reactor, and offered two ways to increase of microflora concentration:

- use of separate cellular agglomerates in the form of "sludge granules" to improve their sedimentation properties and reduce the washing out degree of microflora;
- allow methanogenic bacteria cells to grow in the immobilized form in inert "carriers", which have high specific gravity higher than cells [4, 5].

In this present paper we give results of investigation and evaluation of the performance of single-stage leach-bed reactor for treatment of solid cattle manure, concentrating on the biogas production and biogas composition.


Materials and methods

Feedstock. Cattle manure was provided by livestock farm of the University of Hohenheim. For each experiments and runs were required 3 kg of cattle manure and 19 L of inoculum. The inoculum was fermented cattle slurry that was used as leachate for the microbial initiation of single-phase leach-bed process with immobilization of microorganisms and for recirculation system. Fresh manure (FM) as received and inoculum samples were analyzed in triplicate for its Dry Matter (DM) and moisture content, Organic Dry Matter (oDM) and ash content according to standard methods of APHA (1995) [6]. Accordance with results of tested substrate analysis, DM content of cattle manure was 25.93% and 84.07% of solids were oDM.

Experimental set-up and operation. Bioreactor with a working volume of 50 L had been constructed by modifying leach-bed reactor and fixed-bed reactor. Full characteristics of the experimental set-up are given in Korazbekova et al. (2013) [1]. The experiments were conducted in the biogas laboratory of the State Institute of Agricultural Engineering and Bioenergy of the University of Hohenheim (Stuttgart, Germany). The reactors were maintained at $40 \pm 0,2^{\circ}\text{C}$ and operated at a batch mode until biogas production was detected. Anaerobic digestion was performed in two runs, lasted 28 days for Run 1 and 21 days for Run 2 of Hydraulic Retention Time (HRT) in three replicate indicated as Experiments.

Results and discussion

Experimental reactors were tested during a period of 28 days in Run 1 and 21 days in Run 2 to assess the dry fermentation of cattle manure on a modified leach-bed with immobilization device reactors. Biogas production was depicted by biogas volume and methane content. Cumulative biogas and methane production were determined by summing daily biogas and methane yield, respectively. All the experimental repetitions of 2 runs showed similar findings. To describe the features of the cumulative biogas and methane yield was selected Experiment 1. Results of cumulative biogas production from Experiment 1 are shown in Figure. The rapid initial biogas production was due to readily biodegradable organic matter and presence of high content of the methanogens as depicted in Fig.1. The biogas generation started after inoculating, kept increasing until reaching the peak, and then began to decline (Li et al., 2011) [7]. Run 1 showed the maximum biogas production on day 4 in the amount of $0.032 \text{ Nm}^3 (\text{kg oDM})^{-1}$ and daily biogas generation was observed more than $0.018 \text{ Nm}^3 (\text{kg oDM})^{-1}$ between days 2 and 10, reduced to less than $0.005 \text{ Nm}^3 (\text{kg oDM})^{-1}$ after day 19. The cumulative biogas yield was indicated as $0.327 \text{ Nm}^3 (\text{kg oDM})^{-1}$ at the end of Run 1. Biogas production started faster in Run 2, i.e.



Cumulative biogas production (Experiment 1)

peak biogas generation ($0.047 \text{ Nm}^3 (\text{kg oDM})^{-1}$) was achieved up to day 3 of the experiment demonstrating the maximum degree of biogas formation till days 9 (daily production more $0.02 \text{ Nm}^3 (\text{kg oDM})^{-1}$) and intensive metabolism of microorganisms, since reactor was enriched with methanogens because of biofilm formation and use of fermented leachate from previous run. The cumulative biogas production was $0.328 \text{ Nm}^3 (\text{kg oDM})^{-1}$ at the end of 21-day of HRT. There were not observed several peaks during the digestion process in both two runs as reported by Li et al. (2011) [7].

The curve of cumulative methane production gave volume of $0.148 \text{ Nm}^3 (\text{kg oDM})^{-1}$. Run 2 showed a rapid initiation of methanogenesis from the early days and the maximum methane yield was $0,022 \text{ Nm}^3 (\text{kg oDM})^{-1}$, 1.6 times more than in Run 1 due to the high content of methane-producing bacteria resulted in immobilization of microorganisms on the supporting materials. The total methane yield was $0,150 \text{ Nm}^3 (\text{kg oDM})^{-1}$ at the end of the digestion process.

The biogas quality for Experiment 1. As depicted, the initial percentage of methane in the biogas has increased and exceeded 26.5% after 3 days of start-up, and 50% on day 5 and kept it up to the end of Run 1 presenting stable phase of the digestion. The methane content reached 35% on 2 days and increased sharply to 66% on day 7 as a peak percentage of methane in Run 2. But the maximal methane percentage in Run 1 achieved one day later than in Run 1 on day 8 (56.1%). The percentage of carbon dioxide exceeded the methane percentage during 4 days. The maximum percentage of carbon dioxide was 53.4% on day 2 in Run 1 and 50% on day 3 in Run 2, those were followed by a gradual decrease to 37%. The percentage of carbon dioxide has stabilized in between 42 to 45% from day 6 and 5, respectively in Runs 1 and 2. High initial percentage of carbon dioxide and low methane percentage associated with the biochemical transformation of organic matter in anaerobic digestion. In the second stage of anaerobic digestion (acidogenesis) 70% of low molecular weight compounds, such as simple sugars, amino acids and fatty acids decomposed to acetate, carbon dioxide and hydrogen, and the remaining 30% to volatile fatty acids (VFA) and alcohols [7, 8]. In addition, the regeneration time of acidogenic bacteria (Bacterioids, Clostridia) is 24-36 h [9]. The methane content increased intensively and the maximum percentage also achieved faster for 1 day earlier in Run 2, because Run 2 was initiated with leachate from Run 1, which contained a high concentration of microorganisms [10].

Final data of biogas production on the results of experiments are shown in table. And in accordance with table 1, $(0.331 \pm 0.005) \text{ Nm}^3 (\text{kg oDM})^{-1}$ biogas (standard error = 0.003) with an average percentage of methane $(47.13 \pm 1.40)\%$ was obtained during the anaerobic fermentation of cattle manure in the reactor with immobilization device.

Production and energy content of the biogas from the anaerobic fermentation in the leach-bed reactor

Experiment	Biogas yield, $\text{Nm}^3 (\text{kg oDM})^{-1}$	Average methane content, %	Maximum methane content, %	Methane yield, $\text{Nm}^3 (\text{kg oDM})^{-1}$	Energy value, $\text{kWh} (\text{kg oDM})^{-1}$
Experiment 1 ^a	0.328 ± 0.001	45.50 ± 0.3	61.10 ± 7.10	0.149 ± 0.001	14.8 ± 0.14
Experiment 2 ^a	0.337 ± 0.002	48.10 ± 1.4	68.10 ± 0.14	0.162 ± 0.006	16.1 ± 0.56
Experiment 3 ^a	0.329 ± 0.008	47.75 ± 2.5	65.25 ± 0.07	0.157 ± 0.004	15.6 ± 0.40
Final mean value	0.331	47.13	64.80	0.156	15.50
Standard deviation	0.005	1.40	3.52	0.007	0.66
Standard error	0.003	0.82	2.03	0.004	0.379
Final range ^b	0.331 ± 0.005	47.13 ± 1.40	64.80 ± 3.52	0.156 ± 0.007	15.50 ± 0.66
1 ^a – Data are expressed as mean±standard deviation of two runs.					
2 ^b – Data are expressed as mean±standard deviation of three runs and experiments.					

The maximum methane percentage was found in the value of $(61.10 \pm 7.10)\%$ in Experiment 1, $(68.10 \pm 0.14)\%$ in Experiment 2 and $(65.25 \pm 0.07)\%$ in Experiment 3. The final value was $(64.80 \pm 3.52)\%$ with a standard error of 2.03%. Cumulative methane yield was (0.149 ± 0.001) ; (0.162 ± 0.006) and $(0.157 \pm 0.004) \text{ Nm}^3 (\text{kg oDM})^{-1}$ for Experiments 1, 2 and 3, and the final range was $(0.156 \pm 0.007) \text{ Nm}^3 (\text{kg oDM})^{-1}$ with a standard error of $0.004 \text{ Nm}^3 (\text{kg oDM})^{-1}$. Energy content ranged between 14.8-16.1 in three experiments. Calculated final range lies in the amount of $(15.50 \pm 0.66) \text{ kWh} (\text{kg oDM})^{-1}$ with the standard error of $0.379 \text{ kWh} (\text{kg oDM})^{-1}$.

In conclusion, dry fermentation of cattle manure with dry matter content of 25.93% in leach-bed reactor without anaerobic pretreatment and mixing is feasible and stable process for production of biogas from agricultural wastes. Immobilization of methanogens on the supporting materials (polyethylene packing rings) in leach-bed reactor improved biogas production.

REFERENCES

- [1] Korazbekova K.U., Bakhov Zh.K., Saparbekova A.A. Dry Fermentation of Agricultural Waste in the Modified Leach-bed Reactor with Immobilization of Microorganisms // *Biotechnology*. 2013. Vol. 12(6). P. 236-244.
- [2] Mes T.Z.D., Stams A.J.M., Reith J.H., Zeeman G. Methane production by anaerobic digestion of wastewater and solid wastes. The Netherlands: Dutch Biological Hydrogen Foundation, 2009. Ch. 4. P. 58-102.
- [3] Demirel B., Scherer P. The roles of acetotrophic and hydrogenotrophic methanogens Turing anaerobic conversion of biomass to methane: a review // *Rev. Environ. Sci. Biotechnol.* 2008. N 7. P. 173-190.
- [4] Parawira W. Anaerobic Treatment of agricultural residues and wastewater. Application of high-rate reactors: doctoral dissertation PhD. Sweden, 2004. 53 p.
- [5] Korazbekova K.U., Bakhov Zh.K. Performance of Leach-bed Reactor with Immobilization of Microorganisms in terms of Methane Production Kinetics // *Journal of Biological Sciences*. 2014. Vol. 14(4). P. 236-244.
- [6] APHA. Standard Methods for the Examination of Water and Wastewater. Washington DC, 1995. 53 p.
- [7] Li J., Jha A.K., He J., Ban Q., Chang S., Wang P. Assessment of the effects of dry anaerobic co-digestion of cow dung with waste water sludge on biogas yield and biodegradability // *International Journal of the Physical Sciences*. 2011. Vol. 6(15). P. 3723-3732.
- [8] Seadi T.A., Rutz D., Prassl H., Köttner M., Finsterwalder T., Volk S., Janssen R. Biogas. Handbook. Denmark, 2008. 125 p.
- [9] Deublein D., Steinhauser A. Biogas from Waste and Renewable Resources. Germany, 2008. 423 p.
- [10] Bahov Zh.K., Lemmer A., Korazbekova K.U. Ispol'zovanie reaktora so sloem vyshhelachivaniya i immobilizacii metanogennyh bakterij pri anajerobnom brozhenii navoza // *Izvestija NAN RK. Serija biologicheskaja i medicinskaja*. 2013. N 4(298). P. 155-160.

К. У. Сұлтанғалиева, Ж. К. Бахов, А. А. Абубакирова, Б. Ж. Муталиева

М. Өуезов атындағы Оңтүстік Қазақстан мемлекеттік университеті, Шымкент, Қазақстан

БИОГАЗ ӨНДІРІСІ ҮШІН СІЛТІЛЕНДІРУ ҚАБАТЫ БАР РЕАКТОРДА АУЫЛ ШАРУАШЫЛЫҚ ҚАЛДЫҚТАРЫНЫҢ АНАЭРОБТЫ АШУЫН БАҒАЛАУ

Аннотация. Ауыл шаруашылық қалдықтарының анаэробты ашу процесі экологиялық және әлеуметтік мәселелерді төмендету, сонымен қатар қалпына келтірілетін энергияның альтернативті өндірілуі үшін бүкіл әлемде кең қолданылады. Биогаз өндіруде бір сатылы сілтілендіру қабаты бар процестің тиімділігін бағалау үшін ауыл шаруашылық қалдықтарының (ірі қара мал қиы) құрғақ анаэробты ферментациясы жүргізілді. Тәжірибелер $40 \pm 0,2^\circ\text{C}$ температурада кезеңді режимде жүрді. Жұмыс көлемі 50 л. биореактор сілтілендіру қабаты бар реактор мен бекітілген қабаты бар реакторды модификациялау жолымен құрастырылды. Реактор полиэтилен буылған сақиналар қабаты түріндегі иммобилизациялау қондырғысымен жабдықталған. Ол реактор түбінде орналасқан. Тәжірибелік реакторлар Сынақ 1-де 28 күн, ал Сынақ 2-де 21 күн бойы сыналды. Реактордың өнімділігі биогаз өндіру бойынша талданды. Нәтижелерге сәйкес, анаэробы ашу кезінде биогаздың кумулятивті шығуы $(0,331 \pm 0,005) \text{ Nm}^3(\text{кг оDM})^{-1}$, метанның орташа пайызы $(47,13 \pm 1,40)\%$ құрады. Ауыл шаруашылық қалдықтарының анаэробты ашуы сілтілендіру қабаты бар реакторда араластруды және алдына ала анаэробты өңдеуді қажет етпейтін, жүзеге асыруға болатын және орнықты процесс болып табылады және микрофлораны қондырғыда иммобилизациялау метанның шығуын жақсартады.

Түйін сөздер: анаэробты ашу, биогаз, ауыл шаруашылық қалдықтары, метан, сілтілендіру қабаты бар реактор.

К. У. Султангалиева, Ж. К. Бахов, А. А. Абубакирова, Б. Ж. Муталиева

Южно-Казахстанский государственный университет им. М. Ауэзова, Шымкент, Казахстан

ОЦЕНИВАНИЕ АНАЭРОБНОГО БРОЖЕНИЯ СЕЛЬСКОХОЗЯЙСТВЕННЫХ ОТХОДОВ В РЕАКТОРЕ СО-СЛОЕМ ВЫЩЕЛАЧИВАНИЯ ДЛЯ ПРОИЗВОДСТВА БИОГАЗА

Аннотация. Процессы анаэробного сбраживания сельскохозяйственных отходов широко используются во всем мире для уменьшения экологических и социальных проблем, а также как альтернативное производство возобновляемой энергии. Была проведена анаэробная сухая ферментация сельскохозяйственных отходов (навоз крупного рогатого скота) для оценки эффективности одностадийного процесса со-слоем выщелачивания для производства биогаза. Эксперименты проводились в режиме периодического действия при температуре $40 \pm 0,2^\circ\text{C}$. Биореактор с рабочим объемом 50 л был сконструирован путем модификации реактора выщелачивания и реактора с неподвижным слоем. Реактор был оборудован иммобилизационным устройством, расположенным на дне в виде слоя полиэтиленовых упаковочных колец. Экспериментальные реакторы были испытаны в течение 28 дней в опыте 1 и 21 дней в опыте 2. Производительность реактора была проанализирована с точки зрения производства биогаза. По результатам средний кумулятивный выход биогаза составил $(0,331 \pm 0,005) \text{ Nm}^3 (\text{кг оDM})^{-1}$, средний процент метана $(47,13 \pm 1,40)\%$ в течение анаэробного сбраживания. Анаэробное сбраживание сельскохозяйственных отходов в реакторе со-слоем выщелачивания является осуществимым и стабильным процессом производства биогаза без анаэробной предварительной обработки и смешивания, а иммобилизация микрофлоры на подложке улучшает производство метана.

Ключевые слова: анаэробное сбраживание, биогаз, сельскохозяйственные отходы, метан, реактор со-слоем выщелачивания.

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www.nauka-nanrk.kz

ISSN 2518-1629 (Online), ISSN 2224-5308 (Print)

<http://www.biological-medical.kz/index.php/ru/>

Редактор *М. С. Ахметова, Д. С. Аленов, Т. М. Апендиев*
Верстка на компьютере *Д. Н. Калкабековой*

Подписано в печать 15.05.2017.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
12,4 п.л. Тираж 300. Заказ 3.