

ISSN 2518-1629 (Online),
ISSN 2224-5308 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
Өсімдіктердің биологиясы және биотехнологиясы институтының

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Института биологии и биотехнологии растений

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
of the Institute of Plant Biology and Biotechnology

**БИОЛОГИЯ ЖӘНЕ МЕДИЦИНА
СЕРИЯСЫ**



СЕРИЯ

БИОЛОГИЧЕСКАЯ И МЕДИЦИНСКАЯ



SERIES

OF BIOLOGICAL AND MEDICAL

4 (322)

ШІЛДЕ – ТАМЫЗ 2017 ж.

ИЮЛЬ – АВГУСТ 2017 г.

JULY – AUGUST 2017

1963 ЖЫЛДЫҢ ҚАҢТАР АЙЫНАН ШЫҒА БАСТАҒАН
ИЗДАЕТСЯ С ЯНВАРЯ 1963 ГОДА
PUBLISHED SINCE JANUARY 1963

ЖЫЛЫНА 6 РЕТ ШЫҒАДЫ
ВЫХОДИТ 6 РАЗ В ГОД
PUBLISHED 6 TIMES A YEAR

АЛМАТЫ, ҚР ҰҒА
АЛМАТЫ, НАН РК
ALMATY, NAS RK

Б а с р е д а к т о р

ҚР ҰҒА академигі, м. ғ. д., проф. **Ж. А. Арзықұлов**

Абжанов Архат проф. (Бостон, АҚШ),
Абелев С.К., проф. (Мәскеу, Ресей),
Айтқожина Н.А., проф., академик (Қазақстан)
Акшулаков С.К., проф., академик (Қазақстан)
Алшынбаев М.К., проф., академик (Қазақстан)
Бәтпенев Н.Д., проф., корр.-мүшесі (Қазақстан)
Березин В.Э., проф., корр.-мүшесі (Қазақстан)
Берсімбаев Р.И., проф., академик (Қазақстан)
Беркінбаев С.Ф., проф., (Қазақстан)
Бисенбаев А.К., проф., академик (Қазақстан)
Бишимбаева Н.К., проф., академик (Қазақстан)
Ботабекова Т.К., проф., корр.-мүшесі (Қазақстан)
Bosch Ernesto prof. (Spain)
Жансүгірова Л.Б., б.ғ.к., проф. (Қазақстан)
Ellenbogen Adrian prof. (Tel-Aviv, Israel),
Жамбакин Қ.Ж., проф., академик (Қазақстан), бас ред. орынбасары
Заядан Б.К., проф., корр.-мүшесі (Қазақстан)
Ishchenko Alexander prof. (Villejuif, France)
Исаева Р.Б., проф., (Қазақстан)
Қайдарова Д.Р., проф., академик (Қазақстан)
Кохметова А.М., проф., корр.-мүшесі (Қазақстан)
Күзденбаева Р.С., проф., академик (Қазақстан)
Лось Д.А., prof. (Мәскеу, Ресей)
Lunenfeld Bruno prof. (Израиль)
Макашев Е.К., проф., корр.-мүшесі (Қазақстан)
Муминов Т.А., проф., академик (Қазақстан)
Огарь Н.П., проф., корр.-мүшесі (Қазақстан)
Омаров Р.Т., б.ғ.к., проф., (Қазақстан)
Продеус А.П. проф. (Ресей)
Purton Saul prof. (London, UK)
Рахыпбеков Т.К., проф., корр.-мүшесі (Қазақстан)
Сапарбаев Мұрат проф. (Париж, Франция)
Сарбасов Дос проф. (Хьюстон, АҚШ)
Тұрысбеков Е.К., б.ғ.к., асс.проф. (Қазақстан)
Шарманов А.Т., проф. (АҚШ)

«ҚР ҰҒА Хабарлары. Биология және медициналық сериясы».

ISSN 2518-1629 (Online),

ISSN 2224-5308 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5546-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
www.nauka-nanrk.kz/biological-medical.kz

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2017

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р

академик НАН РК, д.м.н., проф. **Ж. А. Арзыкулов**

Абжанов Архат проф. (Бостон, США),
Абелев С.К. проф. (Москва, Россия),
Айтхожина Н.А. проф., академик (Казахстан)
Акшулаков С.К. проф., академик (Казахстан)
Алчинбаев М.К. проф., академик (Казахстан)
Батпенов Н.Д. проф. член-корр.НАН РК (Казахстан)
Березин В.Э., проф., чл.-корр. (Казахстан)
Берсимбаев Р.И., проф., академик (Казахстан)
Беркинбаев С.Ф. проф. (Казахстан)
Бисенбаев А.К. проф., академик (Казахстан)
Бишимбаева Н.К. проф., академик (Казахстан)
Ботабекова Т.К. проф., чл.-корр. (Казахстан)
Bosch Ernesto prof. (Spain)
Джансугурова Л. Б. к.б.н., проф. (Казахстан)
Ellenbogen Adrian prof. (Tel-Aviv, Israel),
Жамбакин К.Ж. проф., академик (Казахстан), зам. гл. ред.
Заядан Б.К. проф., чл.-корр. (Казахстан)
Ishchenko Alexander, prof. (Villejuif, France)
Исаева Р.Б. проф. (Казахстан)
Кайдарова Д.Р. проф., академик (Казахстан)
Кохметова А.М. проф., чл.-корр. (Казахстан)
Кузденбаева Р.С. проф., академик (Казахстан)
Лось Д.А. prof. (Москва, Россия)
Lunenfeld Bruno prof. (Израиль)
Макашев Е.К. проф., чл.-корр. (Казахстан)
Муминов Т.А. проф., академик (Казахстан)
Огарь Н.П. проф., чл.-корр. (Казахстан)
Омаров Р.Т. к.б.н., проф. (Казахстан)
Продеус А.П. проф. (Россия)
Purton Saul prof. (London, UK)
Рахыпбеков Т.К. проф., чл.-корр. (Казахстан)
Сапарбаев Мурат проф. (Париж, Франция)
Сарбасов Дос проф. (Хьюстон, США)
Турьсыбеков Е. К., к.б.н., асс.проф. (Казахстан)
Шарманов А.Т. проф. (США)

«Известия НАН РК. Серия биологическая и медицинская».

ISSN 2518-1629 (Online),

ISSN 2224-5308 (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов
Министерства культуры и информации Республики Казахстан №5546-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел. 272-13-19, 272-13-18,
www.nauka-nanrk.kz/biological-medical.kz

© Национальная академия наук Республики Казахстан, 2017

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

Editor in chief

Zh.A. Arzykulov, academician of NAS RK, Dr. med., prof.

Abzhanov Arkhat, prof. (Boston, USA),
Abelev S.K., prof. (Moscow, Russia),
Aitkhozhina N.A., prof., academician (Kazakhstan)
Akshulakov S.K., prof., academician (Kazakhstan)
Alchinbayev M.K., prof., academician (Kazakhstan)
Batpenov N.D., prof., corr. member (Kazakhstan)
Berezin V.Ye., prof., corr. member. (Kazakhstan)
Bersimbayev R.I., prof., academician (Kazakhstan)
Berkinbaev S.F., prof. (Kazakhstan)
Bisenbayev A.K., prof., academician (Kazakhstan)
Bishimbayeva N.K., prof., academician (Kazakhstan)
Botabekova T.K., prof., corr. member. (Kazakhstan)
Bosch Ernesto, prof. (Spain)
Dzhansugurova L.B., Cand. biol., prof. (Kazakhstan)
Ellenbogen Adrian, prof. (Tel-Aviv, Israel),
Zhambakin K.Zh., prof., academician (Kazakhstan), deputy editor-in-chief
Ishchenko Alexander, prof. (Villejuif, France)
Isayeva R.B., prof. (Kazakhstan)
Kaydarova D.R., prof., academician (Kazakhstan)
Kokhmetova A., prof., corr. member (Kazakhstan)
Kuzdenbayeva R.S., prof., academician (Kazakhstan)
Los D.A., prof. (Moscow, Russia)
Lunenfeld Bruno, prof. (Israel)
Makashev E.K., prof., corr. member (Kazakhstan)
Muminov T.A., prof., academician (Kazakhstan)
Ogar N.P., prof., corr. member (Kazakhstan)
Omarov R.T., Cand. biol., prof. (Kazakhstan)
Prodeus A.P., prof. (Russia)
Purton Saul, prof. (London, UK)
Rakhypbekov T.K., prof., corr. member. (Kazakhstan)
Saparbayev Murat, prof. (Paris, France)
Sarbassov Dos, prof. (Houston, USA)
Turysbekov E.K., cand. biol., assoc. prof. (Kazakhstan)
Sharmanov A.T., prof. (USA)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of biology and medicine.

ISSN 2518-1629 (Online),

ISSN 2224-5308 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5546-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://nauka-nanrk.kz> / biological-medical.kz

© National Academy of Sciences of the Republic of Kazakhstan, 2017

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF BIOLOGICAL AND MEDICAL

ISSN 2224-5308

Volume 4, Number 322 (2017), 12 – 18

UDC 577.175.14

Z. M. Biyasheva, A. N. Zhumabai, A. M. Shaizadinova, M. Zh. Tleubergenova, S. D. Sarzhanova

Al-Farabi Kazakh National University, Research Institute of Biology and Biotechnology Problems,
Almaty, Kazakhstan.

E-mail: zaremabiya@gmail.com

**MUTAGENIC EFFECTS OF ALPHA RADIATION
IN DROSOPHILA TEST-SYSTEM**

Abstract. There is a long period of time between an agent intervention on a living organism and biological consequences. For this reason, methods for determining a potential mutagenic activity of individual environmental components and natural complexes are required. We used a traditional Muller-5 (Basc) test based on *Drosophila melanogaster* for testing the influence of ecological factors. Due to this test system, we analyzed genetic effects of α -radiation, which is formed during the radioactive decay of radon daughter products. The test system was used to detect mutations with an autonomous manifestation. Now days, it is also used for detecting conditional mutations with non-autonomous manifestation, which form special features related to an invariant part of species appearance of a living organism. The most striking property of conditional mutations is morphoses formation. In our research, the morphoses appeared in the second generation and were due to the presence of conditioned mutations in parents taken from the F_1 . The primary inducer of the conditional mutations emergence was ionizing α -radiation, and in the next generation they were supplemented by the genetic characteristics of the parents being the inversions. The revealed morphoses formed a characteristic group of deformities: blackspots (melanomas) or white spots on a body; curled, curved, or undirected wings; blister on the wings, without one wing, with deformation of the head, thorax and abdomen, mutation of sterility. Sterility was tested in several generations of flies. A characteristic feature of all morphoses is asymmetry and it is defined as a genetically unstable variation of individual morphogenesis associated with changes in the environment. A statistical analysis of experimental data in the Muller-5 test system (Basc) showed that α -radiation has a mutagenic effect with a probability of not less than 95%.

Key words: radon, emanation, α -radiation, inversion, Basc, *Drosophila*, morphoses.

Introduction. Almaty is a city with the highest natural radiation in Kazakhstan, which rich in such natural resources as minerals, metal ores and natural gas and oil reserves. Kazakhstan has 12 % of the world's uranium resources and may be exposed to a variety of hazardous materials including radon, a radioactive gas occurring naturally as an indirect decay product of uranium. Radon gets out of the earth surface through 5 tectonic faults crossing the city territory. Radon and its decay products are sources of α -radiation - a stream of heavy positively charged particles [1]. In nature, alpha particles occur as a result decay of heavy elements atoms, such as uranium, radium and thorium. Emanation (a release of radon into the air pores) happens when the radium decay took place near the soil surface and it was mainly carried out by recoil energy produced by a radon nucleus in the process of radium nucleus disintegration.

Most of radiation is produced not so much from radon but its decay daughter products. Radon emissions are supposed to be dangerous for living organisms and can cause oncological diseases in humans. In human body, radon facilitates some processes also leading to lung cancer. The decay of radon nuclei and its daughter isotopes in the lung tissue causes a micro-burn, as the whole alpha particles energy is absorbed at its decay point. Combination of radon and smoking is especially hazardous and increases the disease risk. According to the US Department of Health, radon had regarded to be the second factor (after smoking) that causes lung cancer, mostly, of bronchogenic (central) type. Lung cancer caused by radon irradiation is the sixth most frequent reason causing death from cancer [2]. Radon

radionuclides cause more than a half radiation dose, which a human body receives from natural and technogenic environmental radionuclides [3-4].

For this reason, the aim of present work is α -radiation mutagenic activity by *Drosophila melanogaster* test-system based on Muller-5 or Basc method.

Materials and methods. The uranium isotope – U^{238} isotope was used as source of α -radiation used. Alpha-rays is one of the ionizing radiation types performing a stream of rapidly moving, positively charged particles (alpha-particles). The main source of this radiation is the radioactive isotopes and daughter products of a natural radon gas. One of the peculiarities of alpha-radiation is its low penetrating power. Alpha particles range in matter (that is a path where ionization is producing) is very short (hundreds of millimeter in biological media, 2.5-8 cm in air). However, along a short path, alpha particles create a great number of ions. That provides a relative biological efficiency, 10 times greater than when exposing the X-ray and gamma radiation.

Testing of α -radiation genetic activity was carried out using the fruit fly *Drosophila melanogaster*. So, some tests based on incidence of different mutations types have been developed for drosophila. The processes occurring in the *Drosophila melanogaster* are extremely interesting for the community of researches engaged in developmental genetics [5]. This fly is chosen as an object in the variety genetic schemes, as it is one of the highly researched and well characterized higher organisms in genetics. Approximately 2/3 of genes that are responsible for a human disease are homologous to genes in *Drosophila melanogaster* genome. The main biochemical processes in *Drosophila melanogaster* and mammalian cells are identical. Also, one of the *Drosophila melanogaster* main advantages lies in the fact that in metabolism process a microsomal activation of substances occurs, when promutagens can be converted into mutagens. This makes it possible to find invisible mutagens, which acquire genotoxicity in metabolism process. Tests based on *Drosophila melanogaster* are recommended by WHO for studying the mutagenic and toxic activity of anthropogenic xenobiotics and pharmacological agents [6].

A traditional Muller-5 (Basc) test based on *Drosophila melanogaster* was used for testing mutagenic activity of α -radiation. Muller-5 method allows to identify lethal and morphological mutations in the F_2 (second generation) X-chromosome. The body of *Drosophila*, like those of all other insects, is divided into segments having certain morphological differences [7]. All flies were identified by their eyes, wings and bristles, because they contained yellow and white genes [8]. We divided males of Oregon wild-type into two samples: the first sample with the males was irradiated by the U^{238} isotope at the exposure of 20-24 hours, and the second control sample was placed nearby, which was not exposed to α -radiation. To obtain F_1 in the Muller-5 (Basc) test system, we used parents different in body and eyes color as well as the shape, because it greatly facilitates females and males identification for parents crossing and second generation analysis.

Presence or absence of males in the population can be determined in a tube without anesthesia. Flies crossing for getting F_1 was carried out massively and individually for F_2 females. The scheme provides an opportunity for F_2 flies crossing without selecting the virgin females. One female from the first generation and two or three M-5 males from the original tube or from the M-5 line have been places in the test tube. For females, such crossing was individual, and the number of tubes was corresponded to half of the analyzed X-chromosomes [9-10]. Sterility was tested using several generations of flies.

The test scheme and a line of flies in the experiment is known as Muller-5 method. This method was developed by H. J. Muller for identifying and recording recessive, sex-linked lethal mutations in drosophila. In the X-chromosome of this line there are 2 inversions – sc^8 and – $sc^{49}(\delta 49)$, which impede a crossing-over between sex chromosomes. The sc^8 inversion captures a major part of X-chromosomes. Since crossing occurs in long inversions, another, shorter inversion $\delta 49$, is introduced into the sc^8 inversion. So, the $\delta 49$ inversion suppresses the cross in the middle region of the X-chromosome. The genes order in $sc^8 \delta 49$ chromosome is violated twice, therefore the cross in it is completely excluded. As a result, both inversions are not associated with a recessive lethal effect, and females homozygous for the Muller-5 chromosome and the same hemizygotic males are viable [6].

A recessive mutation w^a -apricot eyes and dominant mutation Bar-striped eyes serve as phenotypic markers (Figure 1). We obtained two phenotypic classes of females and in the second generation (F_2). In the first generation (F_1) we received B / + females, carrying in the heterozygote X-chromosome of Muller-5 and irradiated male's X-chromosome, and males bearing the Muller-5 X – chromosome in a hemizygot.

The recessive lethal cannot move from the irradiated chromosome to the Muller-5 chromosome, because of locking crossing-over between the irradiated normal chromosome and the Muller-5 chromosome. This makes it possible to bring the irradiated chromosome to a hemizygous state [11].



Figure 1 – Muller-5 line flies: on the left male (smaller) and female on the right

The cultures of the genetic lines and all crosses were kept and propagated on a standard medium [9].

Results and their discussion. Genetic analysis of α -radiation mutagenic activity was carried out according to the scheme shown in Figure 2. Every culture is analyzed visually for revealing morphological mutations after the second-generation flying-out. The conditional mutations are regulatory gene mutations, which are responsible for forming interspecies of similar characteristics [13]. The morphosis formation is a considerable phenomenon of conditional mutations. "Morphosis" means non-inherited morphological disturbances, caused by parent genetic peculiarities. The morphosis, appearing in conditioned mutants, is considered to be different manifestation rates of developmental disturbance. In the case of alpha-radiation induction the offspring were characterized by mutant phenotype of body (white and black plaques, asymmetrical body) and wing: moderate, medium, pronounced, extreme (reduction and even complete disappearance of wing) (Figure 3) [8].




Figure 2 – The scheme of flies crossing according to Muller-5 method for revealing lethal and morphological mutations in X-chromosome: 1 – Muller-5 females; 2 – females with a gray body, with narrow apricot eyes, heterozygote by inducing lethal mutations in X – chromosome; 3 – gray males with narrow apricot eyes; 4 – yellow males with a lethal mutation in X-chromosome are not develop. Irradiated X-chromosome is shown as a bold line [11]

F₂ analysis did not reveal recessive sex-linked mutations. Nevertheless, to evaluate the possible genotoxic effect of α -radiation on individual development of flies, the frequency of occurrence of morphoses was estimated (Table 1).

Table 1– Frequency of morphoses in flies irradiated with α -particles and without it

Sample	Index	Number of flies analyzed (absolute)	Absolute frequency of morphoses	Relative frequency of morphoses, %
With α -radiation (experiment)		3848	28	0.73±0.04
Without α -radiation (control)		3700	10	0.27±0.01

Classical genetics is built on mutations with an autonomous manifestation, based on inheritance signs laws. The Muller-5 (Basc) test system used was designed to evaluate just such mutations. At the present time we began studying mutations with non-autonomous manifestation, which form special features related to an invariant part of species in a living organism appearance. The researchers named such mutations – conditional mutations and received them in *Drosophila* under the influence of ionizing radiation (X-ray irradiation) [13].

One of the main conditions for manifesting a conditional mutation was the presence of chromosome rearrangement in the genotype. At the time being, a conditional mutation is called a local DNA damage, its manifestation being depending on the structure of other genome regions. The most striking property of conditional mutations is the morphoses formation. The term "morphosis" is used to determine non-inherited morphological disorders (deformities) caused by the exposure of extreme environmental factors.

In the genetic literature, morphosis is defined as a non-adaptive and unstable variation of individual morphogenesis, associated with the changes in the external environment. In our experiment, the morphoses were manifested in the second generation (F_2) and were due to the presence of conditioned mutations in parents taken from F_1 . A primary inducer of conditional mutations emergence was α -radiation, and in the second generation they were joined by genetic characteristics of the parents - these are inversions. The morphoses in our experiment formed a very specific group of deformities (Figure 3).




Figure 3 – Morphosis of the second generation according to Muller-5 test-system:

A, B – black plaques on a body; C – moderate mutant wing phenotype (improperly outspread wing);

D – pronounced mutant wing phenotype (improperly outspread wing);

E – extreme mutant wing phenotype – a right wing in the form of an unstructured bubble;

F – morphoses combination - extreme mutant wing phenotype (without a wing), deformation of the head, thorax and abdomen

Most morphoses do not prevent flies from hatching out of pupa, existing, mating, and even giving offspring. The researchers also encounter with cases of morphoses forming, but occurs not so often [13].

As seen in Figure 3, a peculiar feature of all morphoses is asymmetry. They can be distributed over all parts of the body and affect the shape of the head, eyes, chest, legs and wings. Dark spots (or melanomas) similar to necrotic spots that contain conglomerates of dark tissue can appear on all parts of the body.

There can be a single or several individuals containing ugliness may arise. Thus, we found up to 6 morphoses in one F₂ tube. All of them had an individual appearance. The morphoses appeared in F₃, but unlike modifications, they did not reveal phenotypic invariance. So a wing morphosis in F₂ could be revealed as melanoma in F₃ and vice versa. You can say that the type of morphosis is not inherited.

Experimental and control comparison of results was carried out by the Yates' chi-squared test (Table 2) [12].

Table 2 – Experiment and control results in the 2x2 table [14]

Experiment	<i>a</i> (the number of flies without mutation and morphoses)	<i>b</i> (the number of flies with mutation and morphoses)	Σ
		3820	
Control	<i>c</i> (the number of flies without mutation and morphoses)	<i>d</i> (the number of flies with mutation and morphoses)	Σ
		3690	
Σ	7510	38	7548

A statistical experimental data processing in the Muller-5 test-system showed that $\chi^2_{\text{exp}} = 6,99$, a $\chi^2_{\text{table}} = 3,84$ at $k=1$ and $P \leq 0,05$. Therefore at $P \leq 0,05$ $\chi^2_{\text{exp}} > \chi^2_{\text{table}}$. For this reason we can affirm that alpha-radiation possesses a mutagenic effect.

Conclusion. Recessive, sex-linked lethal mutations, modifications and morphoses as the main criterion of α -radiation mutagenic effect evaluation in drosophila have been chosen. Classical genetics is based on mutations with an autonomous manifestation and in our case they are recessive lethal. Mutations with non-autonomous manifestation have been studied quite recently. Due to this, it stands to the reason that the genes, which are responsible for such mutations, form special signs. Basically, these are modifications and morphoses that touch on invariable part of organism's morphology. A common method of mutations evaluation based on *Drosophila melanogaster* test-system has been used in the experiment. The RK Committee for the mutagenicity evaluation of pharmacological preparations recommends this test.

According to the results obtained, a statistically significant difference in the incidence of recessive lethal mutations and conditional mutations induced in the X-chromosome of the drosophila's Oregon line males with alpha irradiation and without it has been shown. The nonparametric chi-square test demonstrated that the frequency distribution control is statistically different at 95% probability level in the experiment and control. Thus, mutagenic activity is revealed in drosophila by alpha-rays irradiation.

Acknowledgments. The Ministry of Education and Science of the Republic of Kazakhstan (grant 2554 / GF4) supported this study.

REFERENCES

- [1] Bersimbaev R., Bulgakova O. (2015) *The health effects of radon and uranium on the population of Kazakhstan*, 37:18. DOI: 10.1186/s41021-015-0019-3
- [2] Darby S.,D. Hill, R. Doll (2001) *Radon: A likely carcinogen at all exposures*, 12:1341-1351. DOI: 10.1023/A:1012518223463
- [3] *Istochniki, jeffekty i opasnost' ionizirujushhej radiacii: Doklad Nauchnogo komiteta OON po dejstviju atomnoj radiacii General'noj Assamblee za 1988 g., s prilozhenijami: V 2-h t. T. 1.: Per. s angl. – M.: Mir, 1992. – S. 552 . il. ISBN:5—03—002458—1*
- [4] *Zashhita ot radona-222 v zhilyh zdaniyah i na rabochih mestah. Publikacija MKRZ 65: Per. s angl. – M.: Jenergoatom izdat, 1995. – S. 68. ISBN: no*

- [5] Marilovtseva E.V., Omelyanchuk L.V. (2015) *hrsGene and Borders of Compartments of Imaginal Wing Disc in Drosophila melanogaster*, DOI: 10.1134/S1022795415100117
- [6] Ashby J. (1994) *International Commission for protection against Environmental Mutagens and Carcinogens. Two million rodent carcinogens. The role of SAR and QSAR in their detection // Mutation Research. vol. 305 (1). – P. 3-12. DOI: no*
- [7] Kyrchanova O.V., Leman D.V., Toshchakov S.V., Utkina M.V., Tikhonov M.V., Parshikov A.F., Maksimenko O.G., Georgiev P.G. (2016) *Induction of Transcription through the scs Insulator Leads to Abnormal Development of Drosophila melanogaster*, DOI: 10.1134/1022795416100057
- [8] Kyrchanova O.V., Georgiev P.G. (2015) *The bithorax Complex of Drosophila Melanogaster as a Model for Studying Specific Long-Distance Interactions between Enhancers and Promoters*, DOI: 10.1134/S1022795415050038
- [9] Bochkov N.P. (2004) *Klinicheskaja genetika*, 475. ISBN: 5-9231-0226-9
- [10] Durnev A.D. (2011) *Geneticheskaja toksikologija // Vestnik Rossijskoj akademii medicinskih nauk, № 9:35-43. (In Russian)*
- [11] Medvedev N. N. (1968) *Prakticheskaja genetika. – 2-oe izd., isp. i dop. 43 – 45; 96-100. ISBN: 978-5-91827-017-2*
- [12] Tihomirova M.M., (1990) *Geneticheskij analiz: Ucheb. Posobie*, 280. ISBN: 5-288-00423-4
- [13] Chadov B.F., E.V. Chadova, S.A. Kopyl, E.V. Artemova, E.A. Hockina, N.B. Fjodorova. (2004) *Ot genetiki vnutrividovyh otlichij k genetike vnutrividovogo shodstva*, 9:1157-1172. DOI: 10.4172/2329-8936.1000137
- [14] Glotov N.V., Zhivotovskij A.A., Hovanov N.V., Hromov-Borisov N.N. (2005) *Biometrics. L.: LGU. 264. ISBN: 5-93972*

З. М. Бияшева, А. Н. Жумабай, А. М. Шайзадинова, М. Ж. Глеубергенова, С. Д. Саржанова

Әл-Фараби атындағы Қазақ Ұлттық университеті, биология және биотехнология мәселелерінің ҒЗУ,
Алматы, Қазақстан

ДРОЗОФИЛАНЫҢ ТЕСТ-ЖҮЙЕСІНДЕГІ α -СӘУЛЕЛЕНУДІҢ МУТАГЕНДІ ӘСЕРІН ТАЛДАУ

Аннотация. Агенттің тірі ағзаға әсер етуі және биологиялық нәтижесі көріну арасында үлкен уақыт айырмашылығы болады, сондықтан қоршаған ортаның жеке компоненттері, комплекстері секілді потенциалды мутагенді белсенділікті анықтау үшін әдістер қажет. Экологиялық стресс-факторлардың генетикалық активтілігін тексеру үшін біз дәстүрлі Меллер-5 (Basc) тест-жүйесін *Drosophila melanogaster*-ге пайдалана отырып жүргіздік. Радонның ыдырау өнімдері кезіндегі негізгі пайда болатын α -сәулеленудің генетикалық эффектісін талдадық. Қолданған тест-жүйе классикалық генетикада мутациялардың автономды көріну жағдайына анықтауға қолданылған. Қазіргі кезде автономды емес шартты мутациялардың көріну жағдайында қолданатын болды. Шартты мутациялар тірі организмнің түр бейнесінің өзгермейтін ерекше белгілерді қамтамасыз етеді. Шартты мутациялардың анық қасиеті – ол морфоздардың пайда болуы. Біздің зерттеуімізде морфоздар екінші ұрпақта пайда болды және бірінші ұрпақтан алынған аталықтардағы шартты мутациялармен негізделінген. Шартты мутациялардың пайда болуына біріншілік индуктор болып иондаушы α -сәулелер болды. Келесі ұрпақта оларды аталықтардың генетикалық ерекшеліктері – инверсиялары толтырды. Табылған морфоздар кемтарлықтардың сипатталған тобын құрастырды: дене бетіндегі қара (меланомалар), немесе ақ дақтар; оралған, майысқан, немесе жайылмаған қанаттар; қанаттардағы көпіршіктер, бір қанатсыз, бастың, көздің, торақтың және қарынның деформациясы, ұрықсыздықтың мутациясы. Ұрықсыздық шыбындардың бірнеше ұрпақтарда тексерілді. Барлық морфоздардың сипатталған белгісі ассиметрия болды және ол қоршаған ортаның өзгерістерімен байланысты жеке морфогенездің генетикалық тұрақты емес вариациясы болып саналады. Меллер-5 (Basc) тест-жүйе статистикалық талдауы α -сәулеленің 95%-дан кем емес сенімділікпен мутагенді әсерін көрсетті.

Түйін сөздер: радон, эманация, α -сәулелену, инверсия, Basc, дрозофила, морфоздар.

З. М. Бияшева, А. Н. Жумабай, А. М. Шайзадинова, М. Ж. Глеубергенова, С. Д. Саржанова

Казахский национальный университет имени аль-Фараби, НИИ проблем биологии и биотехнологии,
Алматы, Казахстан

АНАЛИЗ МУТАГЕННОГО ЭФФЕКТА α -ИЗЛУЧЕНИЯ В ТЕСТ-СИСТЕМЕ ДРОЗОФИЛЫ

Аннотация. Между воздействием агента на живой организм и проявлением биологических последствий проходит часто большой промежуток времени, поэтому необходимы методики определения потенциальной мутагенной активности как отдельных компонентов окружающей среды, так и комплексов. Для проверки генетических эффектов факторов окружающей среды мы использовали традиционный тест

Меллер-5 (Basc) на *Drosophilamelanogaster*. С помощью этой тест-системы мы проанализировали генетические эффекты α -излучения, которое образуется при радиоактивном распаде дочерних продуктов радона. Данная тест-система в классической генетике использовалась для детекции мутаций с автономным проявлением. В настоящее время ее применяют и для обнаружения условных мутаций с неавтономным проявлением, которые формируют особые признаки, относящиеся к инвариантной части видового облика живого организма. Самое яркое свойство условных мутаций – это образование морфозов. В наших исследованиях морфозы проявились во втором поколении и были обусловлены наличием условных мутаций у родителей, взятых из первого поколения (F_1). Первичным индуктором возникновения условных мутаций являлось ионизирующее α -излучение, а в следующем поколении их дополняли генетические особенности родителей – это инверсии. Обнаруженные морфозы составили характерную группу уродств: черные пятна на теле (или меланомы); закрученные, изогнутые, или нерасправленные крылья; белые пятна на теле, пузыри на крыльях, бездонного крыла, с деформацией головы, глаз, торакса и брюшка, мутации стерильности. Стерильность проверялась в нескольких поколениях мух. Характерной чертой всех морфозов является асимметрия и определяется она как генетически не стабильная вариация индивидуального морфогенеза, связанная с изменениями окружающей среды. Статистический анализ данных эксперимента в тест-системе Меллер-5 (Basc) показала, что α -излучение обладает мутагенным эффектом с вероятностью не менее 95%.

Ключевые слова: радон, эманация, α -излучение, инверсия, Basc, дрозофила, морфозы.

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www.nauka-nanrk.kz

ISSN 2518-1629 (Online), ISSN 2224-5308 (Print)

<http://www.biological-medical.kz/index.php/ru/>

Редактор *М. С. Ахметова, Д. С. Аленов, Т. М. Апендиев*
Верстка на компьютере *Д. Н. Калкабековой*

Подписано в печать 27.07.2017.

Формат 60x881/8. Бумага офсетная. Печать – ризограф.
10,7 п.л. Тираж 300. Заказ 4.