

ISSN 2518-1629 (Online),
ISSN 2224-5308 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
Өсімдіктердің биологиясы және биотехнологиясы институтының

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Института биологии и биотехнологии растений

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
of the Institute of Plant Biology and Biotechnology

**SERIES
OF BIOLOGICAL AND MEDICAL**

1 (331)

JANUARY – FEBRUARY 2019

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Б а с р е д а к т о р

ҚР ҰҒА академигі, м. ғ. д., проф. **Ж. А. Арзықұлов**

Абжанов Архат, проф. (Бостон, АҚШ),
Абелев С.К., проф. (Мәскеу, Ресей),
Айтқожина Н.А., проф., академик (Қазақстан)
Ақшулақов С.К., проф., академик (Қазақстан)
Алшынбаев М.К., проф., академик (Қазақстан)
Бәтпенов Н.Д., проф., корр.-мүшесі (Қазақстан)
Березин В.Э., проф., корр.-мүшесі (Қазақстан)
Берсімбаев Р.И., проф., академик (Қазақстан)
Беркінбаев С.Ф., проф., (Қазақстан)
Бисенбаев А.К., проф., академик (Қазақстан)
Бишимбаева Н.Қ., проф., академик (Қазақстан)
Ботабекова Т.К., проф., корр.-мүшесі (Қазақстан)
Bosch Ernesto, prof. (Spain)
Давлетов Қ.К., ассоц.проф., жауапты хатшы
Жансүгірова Л.Б., б.ғ.к., проф. (Қазақстан)
Ellenbogen Adrian, prof. (Tel-Aviv, Israel),
Жамбакин Қ.Ж., проф., академик (Қазақстан), бас ред. орынбасары
Заядан Б.К., проф., корр.-мүшесі (Қазақстан)
Ishchenko Alexander, prof. (Villejuif, France)
Исаева Р.Б., проф., (Қазақстан)
Қайдарова Д.Р., проф., академик (Қазақстан)
Қохметова А.М., проф., корр.-мүшесі (Қазақстан)
Күзденбаева Р.С., проф., академик (Қазақстан)
Локшин В.Н., проф., корр.-мүшесі (Қазақстан)
Лось Д.А., prof. (Мәскеу, Ресей)
Lunenfeld Bruno, prof. (Израиль)
Макашев Е.К., проф., корр.-мүшесі (Қазақстан)
Миталипов Ш.М., (Америка)
Муминов Т.А., проф., академик (Қазақстан)
Огарь Н.П., проф., корр.-мүшесі (Қазақстан)
Омаров Р.Т., б.ғ.к., проф., (Қазақстан)
Продеус А.П., проф. (Ресей)
Purton Saul, prof. (London, UK)
Рахыпбеков Т.К., проф., корр.-мүшесі (Қазақстан)
Сапарбаев Мұрат, проф. (Париж, Франция)
Сарбасов Дос, проф. (Хьюстон, АҚШ)
Тұрысбеков Е.К., б.ғ.к., асс.проф. (Қазақстан)
Шарманов А.Т., проф. (АҚШ)

«ҚР ҰҒА Хабарлары. Биология және медициналық сериясы».

ISSN 2518-1629 (Online),

ISSN 2224-5308 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5546-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://biological-medical.kz/index.php/en/>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Мұратбаева көш., 75.

Г л а в н ы й р е д а к т о р

академик НАН РК, д.м.н., проф. **Ж. А. Арзыкулов**

Абжанов Архат, проф. (Бостон, США),
Абелев С.К., проф. (Москва, Россия),
Айтхожина Н.А., проф., академик (Казахстан)
Акшулаков С.К., проф., академик (Казахстан)
Алчинбаев М.К., проф., академик (Казахстан)
Батпенов Н.Д., проф. член-корр.НАН РК (Казахстан)
Березин В.Э., проф., чл.-корр. (Казахстан)
Берсимбаев Р.И., проф., академик (Казахстан)
Беркинбаев С.Ф., проф. (Казахстан)
Бисенбаев А.К., проф., академик (Казахстан)
Бишимбаева Н.К., проф., академик (Казахстан)
Ботабекова Т.К., проф., чл.-корр. (Казахстан)
Bosch Ernesto, prof. (Spain)
Давлетов К.К., ассоц. проф., ответственный секретарь
Джансугурова Л. Б., к.б.н., проф. (Казахстан)
Ellenbogen Adrian, prof. (Tel-Aviv, Israel),
Жамбакин К.Ж., проф., академик (Казахстан), зам. гл. ред.
Заядан Б.К., проф., чл.-корр. (Казахстан)
Ishchenko Alexander, prof. (Villejuif, France)
Исаева Р.Б., проф. (Казахстан)
Кайдарова Д.Р., проф., академик (Казахстан)
Кохметова А.М., проф., чл.-корр. (Казахстан)
Кузденбаева Р.С., проф., академик (Казахстан)
Локшин В.Н., проф., чл.-корр. (Казахстан)
Лось Д.А., prof. (Москва, Россия)
Lunenfeld Bruno, prof. (Израиль)
Макашев Е.К., проф., чл.-корр. (Казахстан)
Миталипов Ш.М., (Америка)
Муминов Т.А., проф., академик (Казахстан)
Огарь Н.П., проф., чл.-корр. (Казахстан)
Омаров Р.Т., к.б.н., проф. (Казахстан)
Продеус А.П., проф. (Россия)
Purton Saul, prof. (London, UK)
Рахыпбеков Т.К., проф., чл.-корр. (Казахстан)
Сапарбаев Мурат, проф. (Париж, Франция)
Сарбасов Дос, проф. (Хьюстон, США)
Турысбеков Е. К., к.б.н., асс.проф. (Казахстан)
Шарманов А.Т., проф. (США)

«Известия НАН РК. Серия биологическая и медицинская».

ISSN 2518-1629 (Online),

ISSN 2224-5308 (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5546-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел. 272-13-19, 272-13-18,

www.nauka-nanrk.kz / biological-medical.kz

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

Editor in chief

Zh.A. Arzykulov, academician of NAS RK, Dr. med., prof.

Abzhanov Arkhat, prof. (Boston, USA),
Abelev S.K., prof. (Moscow, Russia),
Aitkhozhina N.A., prof., academician (Kazakhstan)
Akshulakov S.K., prof., academician (Kazakhstan)
Alchinbayev M.K., prof., academician (Kazakhstan)
Batpenov N.D., prof., corr. member (Kazakhstan)
Berezin V.Ye., prof., corr. member. (Kazakhstan)
Bersimbayev R.I., prof., academician (Kazakhstan)
Berkinbaev S.F., prof. (Kazakhstan)
Bisenbayev A.K., prof., academician (Kazakhstan)
Bishimbayeva N.K., prof., academician (Kazakhstan)
Botabekova T.K., prof., corr. member. (Kazakhstan)
Bosch Ernesto, prof. (Spain)
Davletov Kairat, PhD, associate professor, executive Secretary
Dzhansugurova L.B., Cand. biol., prof. (Kazakhstan)
Ellenbogen Adrian, prof. (Tel-Aviv, Israel),
Zhambakin K.Zh., prof., academician (Kazakhstan), deputy editor-in-chief
Ishchenko Alexander, prof. (Villejuif, France)
Isayeva R.B., prof. (Kazakhstan)
Kaydarova D.R., prof., academician (Kazakhstan)
Kokhmetova A., prof., corr. member (Kazakhstan)
Kuzdenbayeva R.S., prof., academician (Kazakhstan)
Lokshin V.N., prof., corr. member (Kazakhstan)
Los D.A., prof. (Moscow, Russia)
Lunefeld Bruno, prof. (Israel)
Makashev E.K., prof., corr. member (Kazakhstan)
Mitalipov Sh.M. (America)
Muminov T.A., prof., academician (Kazakhstan)
Ogar N.P., prof., corr. member (Kazakhstan)
Omarov R.T., cand. biol., prof. (Kazakhstan)
Prodeus A.P., prof. (Russia)
Purton Saul, prof. (London, UK)
Rakhypbekov T.K., prof., corr. member. (Kazakhstan)
Saparbayev Murat, prof. (Paris, France)
Sarbassov Dos, prof. (Houston, USA)
Turysbekov E.K., cand. biol., assoc. prof. (Kazakhstan)
Sharmanov A.T., prof. (USA)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of biology and medicine.

ISSN 2518-1629 (Online),

ISSN 2224-5308 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5546-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/> / biological-medical.kz

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF BIOLOGICAL AND MEDICAL

ISSN 2224-5308

Volume 1, Number 331 (2019), 11 – 20

<https://doi.org/10.32014/2019.251-1629.2>

**A. Kaliyev, Ye. Makhambetov, Ye. Medetov, M. Kulmirzayev,
S. Dusembayev, B. Kunakbayev, Ch. Nurimanov, S. Akshulakov**

JSC “National Centre for Neurosurgery”, Department of vascular and functional neurosurgery,
Astana, Kazakhstan.

E-mail: assylbek789@yahoo.com; yermakh@gmail.com; yerkin.medetov@gmail.com;
marat.kulmirzayev@gmail.com; dr.serik@gmail.com; kunakbaevb@gmail.com;
chingiz_nurimanov@mail.ru; raim@rambler.ru

TREATMENT OF COMPLEX INTERNAL CAROTID ANEURYSMS

Abstract.

Introduction: Nowadays different treatment modalities used in the treatment of complex internal carotid artery aneurysms.

Aims: Retrospectively evaluate the results of different treatment modalities other than direct clipping at single.

Design of the study: Retrospective cohort study.

Methods and Material: Clinical presentations, radiological data and outcomes of 64 patients with complex ICA aneurysms evaluated. 15 patients were male, 49 were female with mean age 53 years old.

Results: Follow up period ranged from 3 to 96 months. Parent artery ligation was performed in 12 cases and direct proximal clip placement in 1 case. Endovascular embolization of the aneurysm by coils was performed in 6 cases. Deployment of flow diverter device done in 13 cases. Combined strategy including preliminary bypass with further parent artery occlusion applied in 23 cases. Endovascular occlusion of the parent vessel by coils was done in 9 cases. Surgical morbidity was 20,3%, mortality 3,1%. Outcomes were evaluated by Modified Rankin Scale.

Conclusion: Precise assessment of collateral cerebral blood flow is an important stage in the preoperative planning. Despite new endovascular techniques, EC-IC bypass technique has very important role in the treatment of complex aneurysm. Trapping the aneurysm is still effective and minimally invasive option in selected cases. Combined team approach, with treatment modalities other than direct clipping for complex aneurysms can minimize postoperative morbidity with good outcomes.

Key-words: complex aneurysms, internal carotid artery, EC-IC bypass, flow diverter devices.

Introduction. Despite new endovascular and surgical treatment options, complex intracranial aneurysms are still a big challenge [1-5]. Complex aneurysms recognized as big or giant size, with broad calcified neck, fusiform shape, intraluminal thrombus, branches arising from the aneurysm, atherosclerotic wall of the aneurysms, absence of collateral blood flow[1, 2, 4, 6-9]. These features make such aneurysms difficult for direct surgical treatment with high level of morbidity and mortality [10-22].

For complex aneurysms, multidisciplinary, neurosurgical and interventional neuroradiology team approach is necessary. The aim of the study was to retrospectively evaluate the efficacy, safety and outcomes by methods other than direct clipping.

Subjects and Methods. We conducted retrospective study following experience at our department, between July 2008 and December 2018 to estimate the results of treatment modalities. The review of medical records found 64 patients who underwent endovascular, surgical and multimodality treatment. We reviewed characteristics of aneurysms, treatment modalities, follow up data and results.

Patient's characteristics. Using the criteria of complex aneurysms (table 1) 64 consecutive patients with 64 aneurysms were included into the study. Among 64 patients 15(23,4%) were male, 49(76,6%) were female. Patients age ranged between 19 and 72, with mean age of 53±11,4(Me±SD) years old. Patient's demographic data, clinical presentation, size and configuration of the aneurysm evaluated respectively (table 2). Aneurysm sac size ranged between 11 to 51 mm, with mean size 30±7,9 mm (Me±SD). Size of the neck ranged between 4 to 30 mm, mean size was 10±5,3mm(Me±SD). In 7 cases

fusiform configuration of the aneurysm was revealed. The most frequent location of the aneurysms was paraclinoid (43,8%) and cavernous (35,9%) and aneurysms in the supraclinoid part were found in 20,3%. We did not find significant difference between sex and aneurysm location. In 10(15,6%) cases patients had a history of aneurysmal intracranial bleeding, in 54(84,4%) cases aneurysms were unruptured. Cranial nerves palsy was the leading symptom in clinical presentation in 24(37,5%) cases, 23(35,9%) patients suffered from headache and visual disturbance was found in 17(26,6%) cases.

Table 1 – Criteria for ICA complex aneurysm

Criteria for complex aneurysm	Features
Size	≥11 mm
Neck	≥4 mm
Shape	Regular/Fusiform
Atherosclerotic changes of parent artery, aneurysm dome and neck	Yes/No
Intraluminal thrombus	Yes/No
Absence of collateral blood flow	Yes/No
Presence of ≥3 of listed factors make aneurysm complex.	

Table 2 – Patient’s characteristics

Number of patients	64
Number of aneurysms	64
Mean age	53
Headache	23
Cranial nerves palsy	24
Visual disturbances	17
Ruptured/ Unruptured	10/54
Aneurysm size giant/large	56/8
Cavernous	25
Paraclinoid	21
Supraclinoid	18

Collateral cerebral blood flow assessed by routine balloon test occlusion. Patients treated by endovascular intervention, parent artery occlusion, trapping the aneurysm by bypass surgery or combination of listed methods. Clinical, radiological and angiographic follow up was available. Follow up period ranged between 3-96 months. Outcomes of the treatment assessed by Modified Rankin Scale.

Treatment strategy. Treatment strategy was based on preoperative clinical, radiological and angiographic examination by our multidisciplinary team. Flow chart of treatment strategy presented in table 3. In our department, neurosurgeons are able to perform both microsurgical and endovascular cases. Preoperative assessment of collateral cerebral blood flow performed by double catheter balloon test occlusion. Patients who did not tolerate 30 minutes temporary parent vessel occlusion with 15 minutes drug induced hypotension selected for EC-IC bypass with further endovascular intervention or parent artery occlusion. Selection for either low or high flow bypass done according to anatomy of collateral vessels. Low flow bypass selected in cases when we had to protect of only one territory, such as middle cerebral artery. If there were no any collateral arteries and necessity to cover two main ACA and MCA territories we performed high flow bypass surgery. In some cases, during the angiography and BTO we observed elevation and compression of A1 or M1 segment by aneurysm dome. These patients also underwent preliminary low flow bypass procedures even if they tolerated BTO. This strategy used because of awareness of aneurysm thrombosis, growth and further compression of collateral arteries that could lead to insufficiency of blood circulation and ischemic complications.

Table 3 – Modified Rankin Scale at the time of discharge.


Modified Rankin Scale	Patients number
0	17
1	32
2	8
3	5
4	0
5	0
6	2
Total	64

Surgical treatment included Hunterian parent artery ligation in 12 cases and direct trapping of the aneurysm by proximal and distal clip placement in 1 fusiform aneurysm case. Endovascular embolization of the aneurysm by coils with balloon or stent assistance was performed in 6 cases. In 13 cases we deployed flow diverter devices. Combined strategy including preliminary bypass surgery with further surgical or endovascular intervention was applied in 23 cases. Endovascular occlusion of the parent vessel by coils was done in 9 cases.

Hunterian ligation performed in aneurysms located in the cavernous and paraclinoid part of the ICA. At our later experience, we started to occlude aneurysms located at the paraclinoid and supraclinoid segments by coils at the origin of the ophthalmic artery. This technique implemented because of risk of remnant filling of the aneurysm through external carotid artery system anastomosis.

Results. At the time of data collection, 54 patients (84%) were available for follow up. Follow up period ranged between 3 to 96 months with mean period of 15 months. We observed significant prevalence of female patients among males, and mean age of male was lower than female. Immediate postoperative exclusion of the aneurysm from the circulation was achieved in 61 cases (95,3%). In two cases there was a remnant filling of the aneurysm through meningeal anastomosis from the external carotid artery. These aneurysms were located in the cavernous part of the ICA and did not require additional treatment. Third patient underwent EC-IC bypass and parent artery ligation for ICA supraclinoid aneurysm. On postoperative DSA there was a tiny retrograde filling of the aneurysm through ophthalmic artery. He suffered subarachnoid hemorrhage after two weeks of discharge. He was readmitted to the hospital and underwent additional endovascular embolization of parent artery and remnant aneurysm through the anterior communicating artery. In the group of endovascular embolization with coils, one case of giant aneurysm required second attempt due to significant recanalization of the sac. Combined strategy using bypass procedures with further surgical or endovascular options were performed in 23 cases. 16 patients underwent single STA-MCA bypass, in 6 cases we have done high flow bypass using radial artery graft, in 1 case we created double STA-MCA bypass. After creating an EC-IC bypass patients undergone parent artery occlusion in 18 cases or deployment of flow diverter in 5 cases. Because of small number of in stent thrombosis during our initial experience with flow diverter stent, we did protective bypasses before stent deployment in cases of unfavorable anatomy, insufficient collateral cerebral blood circulation and potential risk of parent artery occlusion due to stent thrombosis. Bypass patency was evaluated using digital subtraction angiography. Early bypass occlusion, during 7 days after the procedure occurred in 2 cases. The longest follow up of bypass patency is 60 months. Surgical associated permanent morbidity occurred in 13 cases (20,3%), two patients (3,1%) died after endovascular stent assistant embolization with coils due to the in stent thrombosis and severe ischemic stroke. At the time of discharge from the hospital outcomes evaluated by Modified Rankin Scale (table 4). Modified Rankin Scale 0 and 1 observed in 49 cases, score 2 was in 8, in 5 cases patients were discharged with score 3. Two patients had score 6. No new aneurysm formation detected. Among patients who were available for MRI follow up, significant aneurysm size decrease observed in cases with parent artery occlusion.

Table 4 – Flow chart of treatment strategy



Discussion. The aim of the treatment of any aneurysm is a total exclusion from the blood circulation. Treatment modalities for complex intracranial aneurysms located at the ICA are still an object for discussion. Both microsurgical and endovascular techniques faces big challenges during the management of these lesions. There are several causes, which are associated with poor outcomes, such as giant size of the dome and neck of the aneurysm, fusiform or dolichoectatic shape, calcified atherosclerotic neck, branches arising from the aneurysm. Such features make these lesions extremely difficult for direct microsurgical clipping or endovascular embolization. However, we think that more aggressive approach for complex aneurysms must be used due to the poor natural history.

According to our retrospective study, we have concluded the importance of multidisciplinary team approach, including neurosurgical and interventional points. Preoperative precise radiological and angiographic assessment of the aneurysm’s dome and neck configuration, presence of intraluminal thrombus and branches arising from the aneurysm should be performed. Special attention to the cerebral collateral blood flow, especially in cases with potential risk of the parent artery occlusion. BOT is a safe and effective method to make a prognosis about tolerance in case of parent artery occlusion [23]. However, there are some reports of ischemic complications even when patients tolerate the BOT [24-28].

In recent report, we performed BOT in all cases, which were scheduled for potential parent artery occlusion. In case of absence of any collateral blood circulation BOT cancelled immediately and we created high flow bypass. Another reason for declining the BOT was hypoplasia of A1 segment of ACA, compression of the A1 or M1 segment by the sac of the aneurysm. In such cases, we decided to create protective low flow bypass. Our decision based on after the acute abruption of the flow in the parent artery and aneurysm there is an increasing volume of the aneurysm due to the thrombosis, that may cause further compression of the surrounding brain tissue and vessels, which may be the reason of ischemic complications and worsening of neurological signs [29, 30]. We have our own experience when a patient tolerated temporary balloon occlusion of the parent artery with further ICA sacrifice. Postoperatively we observed severe neurologic deficit due to the insufficient collateral circulation, emergency EC-IC bypass was performed and a patient did well after the surgery.

Volume decrease of the aneurysm is an important factor, especially in cases with mass effect causing neurological deficit. Parent artery occlusion is related with higher rate of aneurysm sac shrinkage and improvement of symptoms caused by aneurysm size [31-33]. In our experience, among patients treated with therapeutic parent artery endovascular or Hunterian ligation, we could detect significant aneurysm size reduction.

In the past decades, endovascular occlusion of the aneurysms became the first line option in many centers. Outcomes of endovascular treatment of aneurysms became better due to new instruments and devices. However, in some series endovascular management of complex aneurysms is associated with still high rates of recanalization, morbidity and mortality [34]. Introduction of new flow diverter devices promises improvement of the results of endovascular methods. Deployment of flow diverter stents is associated with low level of complications, higher occlusion rates in cases of regular shape aneurysms [35]. Higher morbidity rates of flow diverter deployment occur in cases of complex, giant aneurysms with wide neck [36-38]. In our initial series we encountered technical issues, such as flow diverter migration, stent thrombosis with further occlusion of the parent artery. These complications led to severe ischemic stroke and death in 2 cases. According to this, in our later cases when we anticipated to deploy flow diverter, we created protective bypass if there are no anterior and posterior communicating arteries.

Since the introduction of EC-IC bypass by Yasargil, treatment options for complex aneurysms obtained new opportunities [39]. Trapping the aneurysm by microvascular anastomosis is commonly used worldwide. Complex anatomy of the aneurysm makes complex aneurysms impossible for clipping or endovascular methods. In such cases, EC-IC bypass with further parent artery occlusion sometimes is the only treatment option. Recent reports demonstrated safety, high occlusion rates and long term bypass patency [40]. Furthermore, described bypasses are used as rescue tool in situations of parent artery occlusion and cerebral blood flow insufficiency [41]. At our center, preliminary EC-IC bypass with parent artery occlusion was performed in 22 cases. Current report shows high rates of the aneurysm occlusion, long term bypass patency, and acceptable level of morbidity. In this group, we did not experience any mortality.

The strategy for parent artery occlusion is still controversial. In cases of ICA sacrifice, when the aneurysm is located in the paraclinoid or supraclinoid part there is a risk of remnant flow through the




Figure 1 – Female, 58 y.o. Giant aneurysm of the cavernous part of ICA. Ligation of the ICA was performed. A, B, C preoperatively. Follow up after 6 months (D, E, F): aneurysm is occluded completely. The size of the aneurysm decreased significantly




Figure 2 – Female, 36 y.o. Giant aneurysm of left paraclinoid ICA (A, B, C). STA-MCA bypass with endovascular occlusion of ICA at the level of ophthalmic artery origin. MRI and DSA after 6 months demonstrates exclusion of the aneurysm, volume decrease and bypass patency (D, E, F)




Figure 3 – Female, 51 y.o. with giant aneurysm of paraclinoid left ICA (A, B, C). High flow bypass with endovascular occlusion of ICA. Follow up after 12 months (Pictures D, E, F): exclusion of the aneurysm, bypass patency and aneurysm volume decrease

branches of external carotid artery. Vessels such as ophthalmic or meningo-hypophyseal trunk sometimes can lead to significant aneurysm filling, with reported cases of rupture and aneurysm growth [42]. Our experience with 15 patients when we performed parent artery occlusion in the cervical part of ICA with or without creation of EC-IC bypass showed remnant filling of the aneurysm through ophthalmic artery only in 3 cases. As described previously one of these patients suffered subarachnoid hemorrhage and underwent another surgery. However, now the mechanism of this hemorrhage is not clear. Of course, some of the paraclinoid aneurysms in our series we could perform clipping as well, but our early and midterm follow up of surgical ligation or endovascular ICA occlusion showed good results and low rates of complications.

Conclusion. Current article summarized data of patients with complex internal carotid artery aneurysms treated at single center by methods other than direct clipping. According to our results, assessment of collateral cerebral blood flow is an important stage in the preoperative planning. Major complications in our study were associated with ischemia due to the insufficiency of cerebral circulation. We conclude that whatever treatment option planned, one should pay precise attention to the potential risk of major vessels occlusion. Even in the era of flow diverters, EC-IC bypass technique has very important role in the complex aneurysm management. Parent artery occlusion is still effective and minimally invasive option, but to estimate the results, we need longer follow up and comparison with other treatment modalities. In cases with flow diverter devices risk of complications is higher in the aneurysms with very wide neck, stenosis and severe tortuosity of parent artery. Combined team approach of surgical and endovascular modalities for complex aneurysms can minimize postoperative morbidity and helps to achieve good outcomes.

Source(s) of support: There is no any source of support .

Presentation at a meeting: No any presentations of a given article were organized before.

Conflict of interests. All authors certify that they have no affiliations with or involvement in any organization or entity with any financial interest (such as honoraria; educational grants; participation in speakers' bureaus; membership, employment, consultancies, stock ownership, or other equity interest; and expert testimony or patent-licensing arrangements), or non-financial interest (such as personal or professional relationships, affiliations, knowledge or beliefs) in the subject matter or materials discussed in this manuscript.

All authors have equal contribution to this article, such as data collection, evaluation of results of treatment, attendance to patient care and surgery.

**А. Калиев, Е. Махамбетов, Е. Медетов, М. Кулмирзаев, С. Дюсембаев,
Б. Қунакбаев, Ч. Нуриманов, С. Акшулаков**

"Ұлттық Нейрохирургия Орталығы" АҚ, Тамырлы және функционалды нейрохирургия бөлімшесі,
Астана, Қазақстан

ІШКІ КҮРЕТАМЫРДЫҢ КҮРДЕЛІ АНЕВРИЗМАЛАРЫН ЕМДЕУ

Түйіндеме.

Кіріспе: қазіргі уақытта ішкі күретамырдың күрделі аневризмаларын емдеудің әртүрлі әдістері қолданылуда.

Мақсаты: бір орталықта орындалған тікелей клипстеуден басқа емдеудің әртүрлі әдістерінің қорытындыларын ретроспективтік бағалау.

Зерттеу дизайны: ретроспективті шоғырламалық зерттеу.

Әдістері мен материалдары. Ішкі күретамырдың күрделі аневризмалары бар 64 пациенттің рентгенологиялық деректері және емдеу нәтижелері, клиникалық суреті талданды. 15 пациент – ер адам, 49 пациент – әйел адам, орташа жасы 53 жасты құрады.

Нәтижелері. Операциядан кейінгі бақылау кезеңі 3 айдан бастап 96 айға дейінгі мерзімді құрады. 12 жағдайда тасымалдаушы артерияны жауып тастау және 1 жағдайда ішкі күретамырдың проксималды тікелей клипстеу орындалды. 6 жағдайда микроспиральдарымен аневризмалардың эндоваскулярлық эмболизациясы орындалды. 13 жағдайда ағынды қайта бағыттаушы стентті орнату орындалды. 23 жағдайда тасымалдаушы артерияны кейіннен окклюзиялаумен экстра-краниалды анастомозды алдын-ала салуды

құрайтын біріктірілген стратегия қолданылды. 9 жағдайда микроспиральдарымен ішкі күретамырлардың эндоваскулярлық окклюзиясы орындалды. Операциядан кейінгі асқынулар 20,3%, өлім-жітім 3,1% құрады. Нәтижелер Рэнкиннің түрлендірілген шкаласы бойынша бағаланды.

Қорытынды. Коллатералдық мидың қанағамын бағалау операция алдындағы жоспарлаудың маңызды кезеңі болып саналады. Жаңа эндоваскулярлық әдістерге қарамастан, күрделі аневризмаларды емдеуде экстра-краниалды анастомоз әдісі маңызды рөлді атқарады. Тасымалдаушы артерияны окклюзия жолымен аневризманы окклюзиялау әлі де тиімді және шағын инвазиялық емшара. Күрделі аневризмаларды клипстеуден басқа біріктірілген топтық тәсіл операциядан кейінгі асқынуларды оң нәтижелі жағдайларға дейін жеткізе алады.

Түйін сөздер: күрделі аневризмалар, ішкі күретамыр артериясы, экстра-интракраниалды анастомоз, ағынды қайта бағыттаушы стент.

**А. Калиев, Е. Махамбетов, Е. Медетов, М. Кулмирзаев, С. Дюсембаев,
Б. Қунакбаев, Ч. Нуриманов, С. Акшулаков**

АО Национальный Центр Нейрохирургии, отделение сосудистой и функциональной нейрохирургии,
Астана, Казахстан

ЛЕЧЕНИЕ СЛОЖНЫХ АНЕВРИЗМ ВНУТРЕННЕЙ СОННОЙ АРТЕРИИ

Аннотация.

Введение: в настоящее время применяются различные методы лечения сложных аневризм внутренней сонной артерии.

Цель: ретроспективно оценить результаты различных методов лечения, кроме прямого клипирования выполненных в одном центре.

Дизайн исследования: ретроспективное когортное исследование.

Методы и материалы. Были проанализированы клиническая картина, рентгенологические данные и результаты лечения 64 пациентов со сложными аневризмами ВСА. 15 пациентов – мужчины, 49 – женщины, средний возраст составил 53 года.

Результаты. Период послеоперационного наблюдения составил от 3 до 96 месяцев. Лигирование несущей артерии было выполнено в 12 случаях и прямое клипирование проксимального отдела ВСА в 1 случае. Эндоваскулярная эмболизация аневризм микроспиральями была выполнена в 6 случаях. Установка потока перенаправляющего стента выполнено в 13 случаях. Комбинированная стратегия, включающая предварительное наложение экстра-интракраниального анастомоза с последующей окклюзией несущей артерии, применена в 23 случаях. Эндоваскулярная окклюзия ВСА микроспиральями выполнена в 9 случаях. Послеоперационные осложнения составили 20,3%, летальность 3,1%. Результаты оценивались по модифицированной шкале Рэнкина.

Заключение. Оценка коллатерального мозгового кровотока является важным этапом предоперационного планирования. Несмотря на новые эндоваскулярные методы, метод экстра-интракраниального анастомоза играет очень важную роль в лечении сложных аневризм. Окклюзия аневризм путем окклюзии несущей артерии все еще эффективна и минимально инвазивная процедура. Комбинированный командный подход, кроме прямого клипирования сложных аневризм, может минимизировать послеоперационные осложнения с благоприятными результатами.

Ключевые слова: сложные аневризм, внутренняя сонная артерия, экстра-интракраниальный анастомоз, поток перенаправляющий стент.

Information about authors:

Kaliyev Assylbek, JSC “National Centre for Neurosurgery”, Department of vascular and functional neurosurgery, Astana, Kazakhstan; assylbek789@yahoo.com

Makhambetov Yerbol, JSC “National Centre for Neurosurgery”, Department of vascular and functional neurosurgery, Astana, Kazakhstan; yermakh@gmail.com; <https://orcid.org/0000-0002-7451-8756>

Medetov Yerkin, JSC “National Centre for Neurosurgery”, Department of vascular and functional neurosurgery, Astana, Kazakhstan; yerkin.medetov@gmail.com; <https://orcid.org/0000-0002-0043-0700>

Kulmirzayev Marat, JSC “National Centre for Neurosurgery”, Department of vascular and functional neurosurgery, Astana, Kazakhstan; marat.kulmirzayev@gmail.com; <https://orcid.org/0000-0003-2678-0511>

Dusembayev Serik, JSC “National Centre for Neurosurgery”, Department of vascular and functional neurosurgery, Astana, Kazakhstan; dr.serik@gmail.com; <https://orcid.org/0000-0002-3044-3406>

Kunakbayev Baurzhan, JSC “National Centre for Neurosurgery”, Department of vascular and functional neurosurgery, Astana, Kazakhstan; kunakbaevb@gmail.com; <https://orcid.org/0000-0002-0489-8319>

Nurimanov Chingiz, JSC “National Centre for Neurosurgery”, Department of vascular and functional neurosurgery, Astana, Kazakhstan; chingiz_nurimanov@mail.ru; <https://orcid.org/0000-0002-8251-7980>

Akshulakov Serik, JSC “National Centre for Neurosurgery”, Department of vascular and functional neurosurgery, Astana, Kazakhstan; raim@rambler.ru

REFERENCES

- [1] Hanel R.A., Spetzler R.F. Surgical Treatment of Complex Intracranial Aneurysms // *Neurosurgery*. 2008. 3. P. 1289-1299.
- [2] Choudhri O., Mukerji N., Steinberg G.K. Combined endovascular and microsurgical management of complex cerebral aneurysms // *Endovascular and Interventional Neurology*. 2013. 4. P. 108.
- [3] Velioglu M., Kizilkilic O., Selcuk H., Kocak B., Tureci E., Islak C., Kocer N. Early and midterm results of complex cerebral aneurysms treated with Silk stent // *Neuroradiology*. 2012. 54. P. 1355-1365.
- [4] Shi X., Qian H., Fang T., Zhang Y., Sun Y., Liu F. Management of complex intracranial aneurysms with bypass surgery: a technique application and experience in 93 patients // *Neurosurg Rev*. 2015. 38. P. 109-120.
- [5] Zhu W., Tian Y.L., Zhou L.F., Song D.L., Xu B., Mao Y. Treatment Strategies for Complex Internal Carotid Artery (ICA) Aneurysms: Direct ICA Sacrifice or Combined with Extracranial-to-Intracranial Bypass // *World Neurosurg*. 2011. 75. P. 476-484.
- [6] Jin S.C., Kwon D.H., Song Y., Kim H.J., Ahn J.S., Kwun B.D. Multimodal Treatment for Complex Intracranial Aneurysms: Clinical Research // *J Korean Neurosurg Soc*. 2008. 44. P. 314-319.
- [7] Andaluz N., Zuccarello M. Treatment Strategies for Complex Intracranial Aneurysms: Review of a 12-Year Experience at the University of Cincinnati // *Skull Base*. 2011. 21. P. 233-242.
- [8] Hoh B.L., Putman C.M., Budzik R.F., Carter B.S., Ogilvy C.S. Combined surgical and endovascular techniques of flowalteration to treat fusiform and complex wide-necked intracranial aneurysms that are unsuitable for clipping or coil embolization // *J. Neurosurg*. 2001. 95. P. 24-35.
- [9] Barrow D.L., Cawley C.M. Surgical management of complex intracranial aneurysms // *Neurol India*. 2004. 52. P. 156-162.
- [10] Sharma B.S., Gupta A., Ahmad F.U., Suri A., Mehta V.S. Surgical management of giant intracranial aneurysms // *ClinNeurolNeurosurg*. 2008 110. P. 674-681.
- [11] Dolenc V. Intracavernous aneurysms / in Kaye A., Black P. (eds.) // *Operative Neurosurgery*. New York, 2000. Vol. 2. P. 156-158.
- [12] Cantore G., Santoro A., Guidetti G., Delfinis C.P., Colonnese C., Passacantilli E. Surgical Treatment of Giant Intracranial Aneurysms: Current Viewpoint // *Neurosurgery*. 2008. 63. P. 279-290.
- [13] Shekhtman O.D., Éliava Sh.Sh., Pilipenko Iu.V., Kheïreddin A.S., Okishev D.N., Barchunov B.V., et.al. Long-Term Results of Treatment of Patients with Large and Giant Intracranial Aneurysms of the Internal Carotid Artery // *ZhVoprNeirokhirIm N NBurdenko*. 2013. 77. P. 21-26.
- [14] Xu B.N., Sun Z.H., Romani R., Jiang J.L., Wu C., Zhou D.B., et.al. Microsurgical management of large and giant paraclinoid aneurysms // *World Neurosurg*. 2010. 73. P. 137-146.
- [15] Hiroyuki N., Yasushi S., Yukihido K., Hideyuki O. Long term outcome of unruptured giant cerebral aneurysms // *Neurol Med Chir. Tokyo*, 2006. 46. P. 379-386.
- [16] Kim L.J., Tariq F., Levitt M., Barber J., Ghodke B., Hallam D.K., et.al. Multimodality Treatment of Complex Unruptured Cavernous and Paraclinoid Aneurysms // *Neurosurgery*. 2014. 74. P. 51-61.
- [17] Li J., Lan Z.G., Liu Y., He M., You C. Large and giant ventral paraclinoid carotid aneurysms: Surgical techniques, complications and outcomes // *Clinical Neurology and Neurosurgery*. 2012. 114. P. 907-913.
- [18] Cantore G., Santoro A., Guidetti G., Delfinis C.P., Colonnese C., Passacantilli E. Surgical treatment of giant intracranial aneurysms: current viewpoint // *Neurosurgery*. 2008. 63. P. 279-289.
- [19] Bae H.J., Yoo D.S., Huh P.W., Lee T.G., Cho K.S., Lee S.B. Endovascular Treatment of the Distal Internal Carotid Artery Large Aneurysm // *CerebrovascEndovascNeurosurg*. 2014. 16. P. 200-208.
- [20] Ye G., Zhang M., Deng K., Chen X., Wang Y. Meta-analysis of the Efficiency and Prognosis of Intracranial Aneurysm Treated with Flow Diverter Devices // *J MolNeurosci*. 2016. 59. P. 158-167.
- [21] Jahromi B.S., Mocco J., Bang J.A., Gologorsky Y., Siddiqui A.H., Horowitz M.B., et.al. Clinical and angiographic outcome after endovascular management of giant intracranial aneurysms // *Neurosurgery*. 2008. 63. P. 662-675.

- [22] Akshulakov S.K., Makhambetov E.T., Kaliyev A.B., et al. Surgery of complex aneurysm of the internal carotid artery regarding collateral blood flow. Review of the literature // *News of the National Academy of Sciences of the Republic of Kazakhstan. Series of biological and medical.* 2016. Vol. 5, N 317. P. 11-18.
- [23] Sorteberg W., Boysen, Bakke S.J., Sorteberg A. Angiographic balloon test occlusion and therapeutic sacrifice of major arteries to the brain // *Neurosurgery.* 2008. 63. P. 651-661.
- [24] Andrews J.C., Valavanis A., Fisch U. Management of the internal carotid artery in surgery of the skull base // *Laryngoscope.* 1989. 99. P. 1224-1229.
- [25] Mathis J.M., Barr J.D., Jungreis C.A., Yonas H., Sekhar L.N., Vincent D., et al. Temporary balloon test occlusion of the internal carotid artery: experience in 500 cases // *AJNR Am J Neuroradiol.* 1995. 16. P. 749-754.
- [26] Vazquez Anon V., Aymard A., Gobin Y.P. et al. Balloon occlusion of the internal carotid artery in 40 cases of giant intracavernous aneurysm: technical aspects, cerebral monitoring and results // *Neuroradiology.* 1992. 34. P. 245-51.
- [27] Dare A.O., Gibbons K.J., Gillihan M.D., et al. Hypotensive endovascular test occlusion of the carotid artery in head and neck cancer // *Neurosurg Focus.* 2003. 14. P. 5.
- [28] Dare A.O., Chaloupka J.C., Putman C.M., et al. Failure of the hypotensive provocative test during temporary balloon test occlusion of the internal carotid artery to predict delayed hemodynamic ischemia after therapeutic carotid occlusion // *SurgNeurol.* 1998. 50. P. 147-155.
- [29] Blanc R., Weill A., Piotin M., Ross I.B., Moret J. Delayed Stroke Secondary to Increasing Mass Effect after Endovascular Treatment of a Giant Aneurysm by Parent Vessel Occlusion // *AJNR Am J Neuroradiol.* 2001. 22. P. 1841-1843.
- [30] Turner R.D., Byrne J.V., Kelly M.E., et al. Delayed visual deficits and monocular blindness after endovascular treatment of large and giant paraophthalmic aneurysms // *Neurosurgery.* 2008. 63. P. 469-475.
- [31] De Gast A.N., Sprengers M.E., van Rooij W.J., et al. Midterm clinical and magnetic resonance imaging follow-up of large and giant carotid artery aneurysms after therapeutic carotid artery occlusion // *Neurosurgery.* 2007. 60. P. 1025-1031.
- [32] Clarençon F., Bonneville F., Boch A.L., et al. Parent artery occlusion is not obsolete in giant aneurysms of the ICA. Experience with very-long-term follow-up // *Neuroradiology.* 2011. 53. P. 973-982.
- [33] Maldaner N., Guhl S., Mielke D., et al. For the Giant Intracranial Aneurysm Study Group. Changes in volume of giant intracranial aneurysms treated by surgical strategies other than direct clipping // *ActaNeurochir.* 2015. 157. P. 1117-1123.
- [34] Chalouhi N., Tjoumakaris S., Gonzalez L.F., et al. Coiling of Large and Giant Aneurysms: Complications and Long-Term Results of 334 Cases // *AJNR Am J Neuroradiol.* 2014. 35. P. 546-552.
- [35] Lylyk P., Miranda C., Ceratto R., et al. Curative endovascular reconstruction of cerebral aneurysms with the Pipeline Embolization Device: the Buenos Aires experience // *Neurosurgery.* 2009. 64. P. 632-643.
- [36] Gentric J.C., Darsaut T.E., Makoyeva A., et al. The Success of Flow Diversion in Large and Giant Sidewall Aneurysms May Depend on the Size of the Defect in the Parent Artery // *AJNR Am J Neuroradiol.* 2014. 35. P. 2119-2124.
- [37] Ye G., Zhang M., Deng L., Chen X., Wang Y. Meta-Analysis of the Efficiency and Prognosis of Intracranial Aneurysm Treated with Flow Diverter Devices // *J MolNeurosci.* 2016. 59. P. 158-167.
- [38] Kaliyev A.B. Endovascular treatment of complex ICA aneurysms. Review of the literature // *Neurosurgery and neurology of Kazakhstan.* 2016. Vol. 1(42). P. 19-23.
- [39] Yasargil M.G. *Microsurgery Applied to Neurosurgery.* Stuttgart: Georg Thieme Verlag, 1969. P. 105-115.
- [40] Lee C.H., Chiu T.L., Tsai S.T., Kuo W.C. Extracranial-intracranial bypass in the treatment of complex or giant internal carotid artery aneurysms // *Tzu Chi Medical Journal.* 2015. 27. P. 113-119.
- [41] Pancucci G., Potts M.B., Rodríguez-Hernández A., Andrade H., Guo L., Lawton M.T. Rescue bypass for revascularization after ischemic complications in the treatment of giant or complex intracranial aneurysms // *World Neurosurgery.* 2015. 83. P. 912-920.
- [42] Nakajima N., Nagahiro S., Satomi J., Tada Y., Nakajima K., Sogabe S., et al. Prevention of Retrograde Blood Flow into Large or Giant Internal Carotid Artery Aneurysms by Endovascular Coil Embolization with High-Flow Bypass: Surgical Technique and Long-Term Results // *World Neurosurgery.* 2015. 83. P. 1127-1134.

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www.nauka-nanrk.kz

ISSN 2518-1629 (Online), ISSN 2224-5308 (Print)

<http://biological-medical.kz/index.php/en/>

Редактор *М. С. Ахметова, Т. М. Апендиев, Д. С. Аленов*
Верстка на компьютере *Д. Н. Калкабековой*

Подписано в печать 13.02.2019.

Формат 60x881/8. Бумага офсетная. Печать – ризограф.
6,4 п.л. Тираж 300. Заказ 1.