

ISSN 2518-1629 (Online),
ISSN 2224-5308 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
Өсімдіктердің биологиясы және биотехнологиясы институтының

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Института биологии и биотехнологии растений

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
of the Institute of Plant Biology and Biotechnology

**SERIES
OF BIOLOGICAL AND MEDICAL**

3 (333)

MAY – JUNE 2019

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Б а с р е д а к т о р

ҚР ҰҒА академигі, м. ғ. д., проф. **Ж. А. Арзықұлов**

Абжанов Архат, проф. (Бостон, АҚШ),
Абелев С.К., проф. (Мәскеу, Ресей),
Айтқожина Н.А., проф., академик (Қазақстан)
Ақшулақов С.К., проф., академик (Қазақстан)
Алшынбаев М.К., проф., академик (Қазақстан)
Бәтпенев Н.Д., проф., корр.-мүшесі (Қазақстан)
Березин В.Э., проф., корр.-мүшесі (Қазақстан)
Берсімбаев Р.И., проф., академик (Қазақстан)
Беркінбаев С.Ф., проф., (Қазақстан)
Бисенбаев А.К., проф., академик (Қазақстан)
Бишимбаева Н.Қ., проф., академик (Қазақстан)
Ботабекова Т.К., проф., корр.-мүшесі (Қазақстан)
Bosch Ernesto, prof. (Spain)
Давлетов Қ.К., ассоц.проф., жауапты хатшы
Жансүгірова Л.Б., б.ғ.к., проф. (Қазақстан)
Ellenbogen Adrian, prof. (Tel-Aviv, Israel),
Жамбакин Қ.Ж., проф., академик (Қазақстан), бас ред. орынбасары
Заядан Б.К., проф., корр.-мүшесі (Қазақстан)
Ishchenko Alexander, prof. (Villejuif, France)
Исаева Р.Б., проф., (Қазақстан)
Қайдарова Д.Р., проф., академик (Қазақстан)
Қохметова А.М., проф., корр.-мүшесі (Қазақстан)
Күзденбаева Р.С., проф., академик (Қазақстан)
Локшин В.Н., проф., корр.-мүшесі (Қазақстан)
Лось Д.А., prof. (Мәскеу, Ресей)
Lunenfeld Bruno, prof. (Израиль)
Макашев Е.К., проф., корр.-мүшесі (Қазақстан)
Миталипов Ш.М., (Америка)
Муминов Т.А., проф., академик (Қазақстан)
Огарь Н.П., проф., корр.-мүшесі (Қазақстан)
Омаров Р.Т., б.ғ.к., проф., (Қазақстан)
Продеус А.П., проф. (Ресей)
Purton Saul, prof. (London, UK)
Рахыпбеков Т.К., проф., корр.-мүшесі (Қазақстан)
Сапарбаев Мұрат, проф. (Париж, Франция)
Сарбасов Дос, проф. (Хьюстон, АҚШ)
Тұрысбеков Е.К., б.ғ.к., асс.проф. (Қазақстан)
Шарманов А.Т., проф. (АҚШ)

«ҚР ҰҒА Хабарлары. Биология және медициналық сериясы».

ISSN 2518-1629 (Online),

ISSN 2224-5308 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5546-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://biological-medical.kz/index.php/en/>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р

академик НАН РК, д.м.н., проф. **Ж. А. Арзыкулов**

Абжанов Архат, проф. (Бостон, США),
Абелев С.К., проф. (Москва, Россия),
Айтхожина Н.А., проф., академик (Казахстан)
Акшулаков С.К., проф., академик (Казахстан)
Алчинбаев М.К., проф., академик (Казахстан)
Батпенов Н.Д., проф. член-корр.НАН РК (Казахстан)
Березин В.Э., проф., чл.-корр. (Казахстан)
Берсимбаев Р.И., проф., академик (Казахстан)
Беркинбаев С.Ф., проф. (Казахстан)
Бисенбаев А.К., проф., академик (Казахстан)
Бишимбаева Н.К., проф., академик (Казахстан)
Ботабекова Т.К., проф., чл.-корр. (Казахстан)
Bosch Ernesto, prof. (Spain)
Давлетов К.К., ассоц. проф., ответственный секретарь
Джансугурова Л. Б., к.б.н., проф. (Казахстан)
Ellenbogen Adrian, prof. (Tel-Aviv, Israel),
Жамбакин К.Ж., проф., академик (Казахстан), зам. гл. ред.
Заядан Б.К., проф., чл.-корр. (Казахстан)
Ishchenko Alexander, prof. (Villejuif, France)
Исаева Р.Б., проф. (Казахстан)
Кайдарова Д.Р., проф., академик (Казахстан)
Кохметова А.М., проф., чл.-корр. (Казахстан)
Кузденбаева Р.С., проф., академик (Казахстан)
Локшин В.Н., проф., чл.-корр. (Казахстан)
Лось Д.А., prof. (Москва, Россия)
Lunenfeld Bruno, prof. (Израиль)
Макашев Е.К., проф., чл.-корр. (Казахстан)
Миталипов Ш.М., (Америка)
Муминов Т.А., проф., академик (Казахстан)
Огарь Н.П., проф., чл.-корр. (Казахстан)
Омаров Р.Т., к.б.н., проф. (Казахстан)
Продеус А.П., проф. (Россия)
Purton Saul, prof. (London, UK)
Рахыпбеков Т.К., проф., чл.-корр. (Казахстан)
Сапарбаев Мурат, проф. (Париж, Франция)
Сарбасов Дос, проф. (Хьюстон, США)
Турысбеков Е. К., к.б.н., асс.проф. (Казахстан)
Шарманов А.Т., проф. (США)

«Известия НАН РК. Серия биологическая и медицинская».

ISSN 2518-1629 (Online),

ISSN 2224-5308 (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан **№5546-Ж**, выданное 01.06.2006 г.

Периодичность: 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел. 272-13-19, 272-13-18,
www.nauka-nanrk.kz / biological-medical.kz

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

Editor in chief

Zh.A. Arzykulov, academician of NAS RK, Dr. med., prof.

Abzhanov Arkhat, prof. (Boston, USA),
Abelev S.K., prof. (Moscow, Russia),
Aitkhozhina N.A., prof., academician (Kazakhstan)
Akshulakov S.K., prof., academician (Kazakhstan)
Alchinbayev M.K., prof., academician (Kazakhstan)
Batpenov N.D., prof., corr. member (Kazakhstan)
Berezin V.Ye., prof., corr. member. (Kazakhstan)
Bersimbayev R.I., prof., academician (Kazakhstan)
Berkinbaev S.F., prof. (Kazakhstan)
Bisenbayev A.K., prof., academician (Kazakhstan)
Bishimbayeva N.K., prof., academician (Kazakhstan)
Botabekova T.K., prof., corr. member. (Kazakhstan)
Bosch Ernesto, prof. (Spain)
Davletov Kairat, PhD, associate professor, executive Secretary
Dzhansugurova L.B., Cand. biol., prof. (Kazakhstan)
Ellenbogen Adrian, prof. (Tel-Aviv, Israel),
Zhambakin K.Zh., prof., academician (Kazakhstan), deputy editor-in-chief
Ishchenko Alexander, prof. (Villejuif, France)
Isayeva R.B., prof. (Kazakhstan)
Kaydarova D.R., prof., academician (Kazakhstan)
Kokhmetova A., prof., corr. member (Kazakhstan)
Kuzdenbayeva R.S., prof., academician (Kazakhstan)
Lokshin V.N., prof., corr. member (Kazakhstan)
Los D.A., prof. (Moscow, Russia)
Lunefeld Bruno, prof. (Israel)
Makashev E.K., prof., corr. member (Kazakhstan)
Mitalipov Sh.M. (America)
Muminov T.A., prof., academician (Kazakhstan)
Ogar N.P., prof., corr. member (Kazakhstan)
Omarov R.T., cand. biol., prof. (Kazakhstan)
Prodeus A.P., prof. (Russia)
Purton Saul, prof. (London, UK)
Rakhypbekov T.K., prof., corr. member. (Kazakhstan)
Saparbayev Murat, prof. (Paris, France)
Sarbassov Dos, prof. (Houston, USA)
Turysbekov E.K., cand. biol., assoc. prof. (Kazakhstan)
Sharmanov A.T., prof. (USA)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of biology and medicine.

ISSN 2518-1629 (Online),

ISSN 2224-5308 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5546-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/biological-medical.kz>

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF BIOLOGICAL AND MEDICAL

ISSN 2224-5308

Volume 3, Number 333 (2019), 48 – 55

<https://doi.org/10.32014/2019.2519-1629.30>

UDC 633.18

G. Zh. Medeuova

Kazakh National Women Teachers Training University, Almaty, Kazakhstan.

E-mail: medeuova.galiya96@gmail.com

RICE VARIETIES' AGROCOENOSIS FORMATION AND GRAIN CROPPING CAPACITY DEPENDING ON FERTILIZER QUANTITY

Abstract. Plant density in agrocenosis and the number of productive stems have significant effect on formation of a high grain yield of rice varieties and samples differing in height and architectonics. Thus, when an optimum fertilizer dosage (N180P120 kg/ha a.v.) is added, high grain yield is formed under the following conditions: 1) a high grain yield of a medium growth variety with narrow, vertically arranged leaves (Kuban 3, Krasnodar 424 and Dubovsky 129 varieties) is formed for the density of rice standing on the shoots of 320-400 pcs/m², before harvesting - 250-350 pcs/m², with a feed area of one plant of 22-30 cm², in the period of sweeping - 25-38 cm². In such an agrocenosis, the number of productive stems is 550-650 pcs/m², or 5.5-6.5 million panicles per hectare. 2) A high grain yield in medium-sized, large-leaved varieties (Marzhan, Aral 202, Togusken 1) is formed on highly productive crops, with a standing density of 280-350 pcs/m², before harvesting - 220-250 pcs/m², with 460-580 pcs./m² of productive stalks, or 4.6-5.8 million panicles per hectare. 3) In stunted, with large and wide leaves of varieties (Estuary) and rice samples (ECM), high grain yield is formed with a standing density of 280-380 pcs/m², before harvesting 220-310 pcs/m², with formations 550-650 pcs/m² of productive stems, or 5.5-6.5 million panicles.

Key words: rice, variety, effect of fertilizers on agrocenosis structure, rice varieties yield dependence on fertilizers' dosages.

In rice agrocenosis, formation of the number of productive stems changes during the growing season and, as an integral indicator, depends on many factors: the number and quality of sown seeds, the number of germinated seedlings, their rooting and preservation, tillering intensity, growth and productivity of side shoots. The number of these indicators, the intensity of growth and development of lateral shoots are significantly influenced by doses and methods of applying mineral fertilizers. The influence of the seeding rate and plant stand density in crops on the yield of rice grain has been studied in many studies and experimental results have been accumulated. In these works, the optimal plant stand density in crops is considered as one of the main factors that have a significant impact on the formation of high yields [1-3]. Currently, in Asian countries they intensify the technology of this valuable culture cultivation in order to provide their population with rice. Thus, according to scientists and specialists from China [4], Indonesia [5], Korea [6] and Vietnam [7], the highest and high-quality grains of rice are formed on crops with optimal plant density, balanced nutrients, and optimal doses and methods of mineral fertilizer application and with timely and high-quality carrying out of all agrotechnical techniques. The influence of the seeding rate and plant stand density on the yield of rice grain has been studied in many studies and experimental results have been accumulated. In these works, the optimum plant stand density is considered as one of the main factors that have a significant impact on the formation of high yields [1-3, 8].

Methods and objects of research. Small-plot and large-field field experiments were carried out on the saline soils of the Kazakhstan Aral Sea area in 2012-2017 to determine the favorable structure of agrocenosis of rice varieties and the optimum dosage of nitrogen fertilizer. Medium-height, narrow-leaved, early-ripe varieties such as Kuban 6 and Dubovsky 129, middle-ripe Krasnodar 424 variety, medium-tall, large-leaved samples 4-15, Kyzylorda 5, KzROS 356, Marzhan variety, low-growing, broad-leaved Firth were studied. In small-scale experiments, the seeding rate was 100, 300, 500, 700, 900 kg/m²;

the dosage of fertilizers N90P90, N180P120 kg/ha a.v., were repeated three times. In large-scale experiments, the seeding rate for Kuban 3 and Liman varieties was 100 kg/ha (3 million viable seeds per hectare), 200 kg/ha (6 million), 250 kg/ha (7.5 million), 300 kg/ha (9.5 million viable seeds per hectare). Seeds of Marzhan variety are larger and spinous, therefore they sowed 130 kg/ha (3 million viable seeds), 230 kg/ha (6 million), 280 kg/ha (7.5 million) and 330 kg/ha (9.5 million viable seeds) using a seeder. Pilot crops were flooded on May 18-20. Mineral fertilizers (ammonium sulfate, granulated superphosphate) were added: N0P0, N90P90, N120P120, N150P120, N180P120, N240P120 kg/ha ae. The area of the plots was 150 m², the number of replications - four.

Results of research and discussion. *The main indicators of the structure of rice agroecosystem* include the optimum plant standing density, the number and productivity of main and lateral shoots and tillering intensity. These indicators vary significantly depending on the area of nutrition and doses and methods of mineral fertilizer application. Thus, in model experiments (plot area - 5 m²), the greatest germination of seeds was observed when 100, 300 seeds were sown per m². With an increase in the number of sown seeds (i.e. with a decrease in the area of nutrition), field germination of seeds decreased. For example, the germination of seeds of narrow-leaved rice varieties Kuban 3, Krasnodarskiy 424 and Dubovskiy 129 decreased from 76.0-79.3% to 48.2-57.4%. The same patterns were observed with crops of large-leaved varieties Marzhan and samples 4-15, Kyzylorda 5, KSROC 356. However, with an increase in the seeding rate, more sprouting rice plants and the productive stems remained till harvesting (Fig. 1). Rice is an intensely bushy plant. With increase in area of nutrition and increase of fertilizers dosages, the number of the 2nd, 3rd and other side shoots increased significantly. However, with a decrease in the area of nutrition and the emergence of optimally dense shoots, tillering of rice decreases, shoots of the first and second order appear, which grow and develop synchronously, not lagging behind the main stem. This is one of the main conditions for obtaining high grain yields. But, with very thickened crops, the leaves located on the lower and middle levels are mutually strongly shadowed, tillering lingers, shoots appear without panicle, which leads to a decrease in grain yield [1-3].

When an optimum fertilizer dosage is added (N180/P120 kg/ha) and with highly productive agroecosystem (i.e. at the optimally crowded cultures) tillering increases and the number of productive stalks and side shoots (stalks with heads) (figure). Thus, when 300, 500, 700 pcs of germinating seeds per a m² and adding fertiliser at the rate of N180P120 kg/ha number of productive stalks of the pilot sowings of large-leaf varieties was 452-567, 469-613, 479-634 pcs/m², correspondingly. As for the narrow-leaf rice varieties with vertical leaf position, such as Kuban 3, Krasnodarskiy 424, the number of productive stalks was 487-686, 569-667 pcs/m², i.e. 8.8-21.0% more. At the optimum fertilizer dosage, such an increase in the number of productive stalks is the main condition of high grain yield formation (figure).

When an optimum fertilizer dosage is added to the rice varieties with narrow vertically arranged leaves such as Kuban 3 and Krasnodarskiy 424, the best yield was achieved when 500, 700 pcs/m² of germinating seeds had been sown and at the density of plants growing of 320-430 pcs/m², prior to harvest – 250-280 pcs./m², at the area of one plant feed of 22-30 cm, during the period of ear formation – 25-38 cm², at the number of productive stalks of 550-650 pcs/m², i.e. when 5.5-6.5 mln. heads are formed per a hectare.

As regards the crops of large-leaved varieties and samples of rice, a high grain yield is formed when sowing 300, 500 pcs / m² of viable seeds and with formations of plant standing on shoots of 230-350 pcs/m², in the period of sprouting 220-300 pcs / m², with one feed area plants shoots 28-45 cm², in the period of sweeping - 31-50 cm². Under such conditions, the optimal number of productive stems was 460-580 pcs / m², i.e. 4.6-5.8 million ears of corn were formed on one hectare. Such crops are called by us highly productive agroecosystem [1-3].

Rice crops - dynamic agroecosystem. With the introduction of the optimal dose of fertilizers and depending on the density of standing and feeding area, the mutual influence of neighboring rice plants in crops during the growing season changes. These changes depend on the architectonics, morphophysiological and genetic characteristics of rice varieties. In rice crops, photosynthetic activity and plant growth processes have a significant impact on the level of yield and grain quality depending on standing density. In this regard, the rice crops are subdivided as follows: thinned, moderate, highly productive and highly thickened agroecosystem. The yield of such crops vary [1-3].




Figure 1 – Formation of agroecological structure of rice medium-growth varieties and samples differing by the architectonic depending on fertilisers' dosages and coverage area.

Conventional designation: 1 – number of springing rice sprouts, pcs./m²; 2 – number of plants before harvesting, pcs./m²; 3 – number of productive rice stalks at the fertilizer dosage of N90P90kg/ha; 4 – number of productive rice stalks at the dosage of N180P120 kg/ha; 5 – field germination rate %; 6 – rice tillering capacity at the fertilizer dosage of N90P90 kg/ha; 7 – rice tillering capacity at the fertilizer dosage of N180P120 kg/ha.

On all the different soils of the CIS territories, rice is responsive to the doses and methods of applying nitrogen fertilizers compared to phosphate fertilizers. However, with the joint application of nitrogen and phosphate fertilizers, rice forms the highest grain yield [1-3]. In this regard, the effect of nutrition area and doses of nitrogen-phosphate fertilizers on the formation of the structure and yield of agroecological structure of rice varieties (table) was studied.

Germinating ability of seeds and survival ability of sprouts. There is a positive correlative relationship between field seed germination, the number of preserved plants, the density of standing productive stalks and grain yield of rice crops [1-3]. Thus, when sowing 100, 300 seeds per a m² within the framework of the on model experiments (plot area 5 m²), field seed germination was high. With an increase in the number of seeds sown (i.e. with a decrease in the nutrition area), field germination decreased. For example, field germination of seed varieties Kuban 3, Krasnodar 424, Marzhan decreased from 74-82% to 51.3-53.7%. Despite this, with an increase in the seeding rate, seedlings sprouted thick, the number of plants on crops increased and remained until harvest. Such patterns were preserved on experimental crops of dwarf and low-growing rice samples (figure; table). In model experiments, when sowing 100, 300, 500, 700 pcs / m² of viable seeds and when making optimal doses of fertilizers, the survival rate of the seedlings was high and more rice plants remained. But when sowing 900 pieces / m² of seed, the survival rate of seedlings decreased significantly. The results of the experiments show that with

Formation of rice density in agrocenosis and grain yield depending on increasing dosages of nitrogen fertilizers and seeding rate

| Rice Variety | Seeding Rate, kg/ha | Number of Plants pcs/m ² | | Field Germination Capacity, % | Sprouts Survival Rate, % | Number of Productive Stalks, pcs/m ² | Tilling Capacity, pcs/plant | Grain Yield, h.kg/ha |
|-------------------|---------------------|-------------------------------------|----------------|-------------------------------|--------------------------|---|-----------------------------|----------------------|
| | | Sprouts | Before Harvest | | | | | |
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| N0P0, Control | | | | | | | | |
| Kuban 3 | 100 | 144 | 110 | 48,0 | 76,4 | 186 | 1,69 | 33,4 |
| | 200 | 243 | 189 | 40,5 | 77,8 | 254 | 1,34 | 37,0 |
| | 250 | 300 | 240 | 40,0 | 80,0 | 301 | 1,25 | 38,1 |
| | 300 | 370 | 286 | 38,9 | 77,4 | 334 | 1,17 | 41,8 |
| HCP ₀₅ | – | 13,2 | 14,1 | – | – | 12,6 | – | 4,0 |
| N120P120 kg/ha | | | | | | | | |
| Kuban 3 | 100 | 151 | 116 | 50,3 | 76,8 | 480 | 4,22 | 42,2 |
| | 200 | 243 | 194 | 40,5 | 79,8 | 513 | 2,64 | 51,5 |
| | 250 | 314 | 247 | 41,9 | 78,7 | 531 | 2,15 | 58,1 |
| | 300 | 370 | 287 | 38,9 | 77,6 | 519 | 1,81 | 55,5 |
| HCP ₀₅ | – | 15,3 | 13,4 | – | – | 16,7 | – | 4,3 |
| N150P120 kg/ha | | | | | | | | |
| Kuban 3 | 100 | 143 | 109 | 47,7 | 76,2 | 508 | 4,66 | 50,6 |
| | 200 | 237 | 187 | 39,8 | 78,2 | 518 | 2,72 | 54,1 |
| | 250 | 305 | 243 | 40,5 | 79,7 | 553 | 2,28 | 67,7 |
| | 300 | 366 | 291 | 38,5 | 79,5 | 534 | 1,84 | 60,4 |
| HCP ₀₅ | – | 14,8 | 14,2 | – | – | 16,0 | – | 3,3 |
| N180P120 kg/ha | | | | | | | | |
| Kuban 3 | 100 | 148 | 112 | 49,3 | 75,7 | 549 | 4,90 | 60,8 |
| | 200 | 250 | 200 | 41,7 | 80,0 | 571 | 2,86 | 66,3 |
| | 250 | 319 | 253 | 42,7 | 79,3 | 591 | 2,34 | 78,0 |
| | 300 | 380 | 295 | 40,0 | 77,6 | 658 | 2,23 | 65,9 |
| HCP ₀₅ | – | 15,5 | 13,7 | – | – | 14,2 | – | 4,6 |
| N240P120 kg/ha | | | | | | | | |
| Kuban 3 | 100 | 146 | 116 | 48,7 | 79,5 | 581 | 5,01 | 66,2 |
| | 200 | 248 | 196 | 41,3 | 79,0 | 600 | 3,06 | 73,2 |
| | 250 | 315 | 245 | 42,0 | 77,8 | 631 | 2,58 | 74,0 |
| | 300 | 376 | 298 | 39,6 | 79,3 | 672 | 2,26 | 58,8 |
| HCP ₀₅ | – | 13,5 | 12,8 | – | – | 13,8 | – | 4,5 |
| N0P0, control | | | | | | | | |
| Mar-zhan | 130 | 124 | 95 | 41,3 | 76,6 | 176 | 1,85 | 35,3 |
| | 230 | 224 | 175 | 37,3 | 78,1 | 235 | 1,34 | 38,7 |
| | 280 | 278 | 219 | 38,1 | 78,8 | 278 | 1,27 | 41,3 |
| | 330 | 336 | 273 | 35,4 | 82,3 | 273 | 1,16 | 42,4 |
| HCP ₀₅ | – | 12,1 | 13,4 | – | – | 10,8 | – | 4,3 |
| N120P120 kg/ha | | | | | | | | |
| Mar-zhan | 130 | 123 | 102 | 41,2 | 82,8 | 505 | 4,95 | 44,8 |
| | 230 | 227 | 175 | 37,8 | 77,1 | 512 | 2,93 | 53,4 |
| | 280 | 296 | 227 | 39,5 | 76,7 | 535 | 2,36 | 61,5 |
| | 330 | 345 | 268 | 36,3 | 77,7 | 518 | 1,93 | 51,2 |
| HCP ₀₅ | – | 12,3 | 10,3 | – | – | 12,0 | – | 4,6 |

| Continuation of table | | | | | | | | |
|-----------------------|-----|------|------|------|------|------|------|------|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| N150P120 kg/ha | | | | | | | | |
| Mar-zhan | 130 | 119 | 103 | 39,7 | 86,6 | 516 | 5,01 | 51,5 |
| | 230 | 228 | 179 | 38,0 | 78,5 | 528 | 2,95 | 57,4 |
| | 280 | 293 | 231 | 39,1 | 78,8 | 549 | 2,38 | 70,2 |
| | 330 | 349 | 276 | 36,7 | 79,1 | 569 | 2,05 | 57,8 |
| HCP ₀₅ | – | 13,7 | 12,0 | – | – | 14,0 | – | 4,4 |
| N180P120 kg/ha | | | | | | | | |
| Mar-zhan | 130 | 125 | 98 | 41,7 | 78,4 | 522 | 5,33 | 59,0 |
| | 230 | 233 | 173 | 38,8 | 74,2 | 541 | 3,13 | 66,8 |
| | 280 | 297 | 229 | 39,6 | 77,1 | 551 | 2,41 | 73,8 |
| | 330 | 357 | 282 | 37,6 | 79,0 | 591 | 2,10 | 61,5 |
| HCP ₀₅ | – | 14,1 | 12,7 | – | – | 13,5 | – | 4,5 |
| N240P120 kg/ha | | | | | | | | |
| Mar-zhan | 130 | 121 | 97 | 40,7 | 80,2 | 539 | 5,56 | 60,1 |
| | 230 | 229 | 171 | 38,2 | 74,7 | 557 | 3,26 | 66,7 |
| | 280 | 291 | 225 | 38,8 | 77,3 | 594 | 2,64 | 65,5 |
| | 330 | 361 | 289 | 38,0 | 80,1 | 624 | 2,16 | 53,0 |
| HCP ₀₅ | – | 5,3 | 5,5 | – | – | 6,4 | – | 3,8 |
| N0P0, Control | | | | | | | | |
| Liman | 100 | 121 | 94 | 40,3 | 77,3 | 163 | 1,73 | 28,8 |
| | 200 | 222 | 163 | 37,0 | 73,4 | 228 | 1,40 | 32,7 |
| | 250 | 269 | 210 | 35,9 | 78,1 | 263 | 1,25 | 35,0 |
| | 300 | 335 | 252 | 35,3 | 75,2 | 314 | 1,25 | 38,4 |
| HCP ₀₅ | – | 12,5 | 12,8 | – | – | 11,4 | – | 4,6 |
| N120P120 kg/ha | | | | | | | | |
| Liman | 100 | 119 | 100 | 39,7 | 84,0 | 490 | 4,90 | 42,4 |
| | 200 | 219 | 167 | 36,5 | 76,3 | 514 | 3,08 | 57,5 |
| | 250 | 284 | 217 | 37,9 | 76,4 | 530 | 2,44 | 64,2 |
| | 300 | 341 | 257 | 35,9 | 75,4 | 518 | 2,02 | 57,4 |
| HCP | – | 14,2 | 13,1 | – | – | 12,3 | – | 5,0 |
| N150P120 kg/ha | | | | | | | | |
| Liman | 100 | 124 | 96 | 41,3 | 77,4 | 515 | 5,36 | 51,9 |
| | 200 | 224 | 174 | 37,3 | 77,7 | 558 | 3,21 | 65,0 |
| | 250 | 287 | 232 | 38,3 | 80,8 | 601 | 2,60 | 71,6 |
| | 300 | 347 | 265 | 36,5 | 76,4 | 540 | 2,04 | 61,9 |
| HCP ₀₅ | – | 14,4 | 15,2 | – | – | 13,7 | – | 4,1 |
| N180P120 kg/ha | | | | | | | | |
| Liman | 100 | 123 | 95 | 41,2 | 77,4 | 553 | 5,76 | 57,0 |
| | 200 | 228 | 167 | 38,0 | 73,2 | 571 | 3,42 | 66,5 |
| | 250 | 286 | 228 | 38,1 | 79,7 | 641 | 2,81 | 69,2 |
| | 300 | 352 | 273 | 37,1 | 77,6 | 675 | 2,47 | 57,1 |
| HCP ₀₅ | – | 15,5 | 14,3 | – | – | 13,1 | – | 4,8 |
| N240P120 kg/ha | | | | | | | | |
| Liman | 100 | 123 | 98 | 41,0 | 79,7 | 581 | 5,93 | 52,7 |
| | 200 | 227 | 160 | 37,8 | 70,5 | 600 | 3,75 | 58,5 |
| | 250 | 285 | 227 | 38,0 | 79,6 | 659 | 2,90 | 58,2 |
| | 300 | 345 | 275 | 36,4 | 79,5 | 691 | 2,51 | 53,2 |
| HCP ₀₅ | – | 13,4 | 12,3 | – | – | 15,0 | – | 3,7 |

the introduction of the optimal dose of fertilizers (N180P120 kg / ha ai) and with the optimum density of germinated seedlings, the penetration of light, carbon dioxide (CO₂) and the temperature inside the agroecosystem improved and good conditions were established for the growth and development of each rice plant in agroecosystem.

Rice tillering depends on the area of nutrition [1-3], the dose, timing and methods of applying mineral fertilizers [1-3, 9, 10, 16], the depth of water within the check plots and changes in irrigation mode [9, 10, 12, 13, 15]. The results of our research showed that with an increase in the seeding rate (i.e., a decrease in the nutritional area) and an increase in plant density to the optimum level, tillering of rice stops. With the introduction of an average dose (N120P90-120 kg / ha), tillering of rice increases as compared with plants on the control (without fertilizer) and the introduction of small doses (N90P90 kg / ha AI) of fertilizers. With an increase in the dose of fertilizer to the optimum (N180P120 kg / ha) and sowing of 500, 700, 900 pcs / m², the level of tillering of rice decreased. In spite of this, with increasing doses of fertilizers, the number of productive stems increased. On crops of medium-grade rice varieties, with the introduction of optimal (N180P120 kg / ha) and high (N240P180 kg / ha AV) fertilizer dose, bushiness when sown 100 pcs / m² from 7.7-9.1 pcs / m² decreased to 1, 8-1.9 pcs / m², i.e. 4.1-4.7 times when sown 900 pieces / m² of germinating seeds. The same patterns were observed on dwarf and low-growing varieties. However, with increasing doses of fertilizers, bushiness and the number of productive rice stalks increased to a certain extent (figure). Consequently, on high agro-backgrounds, a decrease in nutritional area (i.e., an increase in plant standing density) is a factor that reduces rice bushiness.

The productivity of lateral shoots from the different rows is not the same. As the place of emergence of lateral shoots in the tillering node increases, the stem height, the length of the panicle and the number of grains on the panicle of side shoots decrease. Side shoots that appear on the lower level of the tillering node, i.e. 3-5 leaves on the bosom grows and develops synchronously, not lagging behind the main stem and is the most productive. However, depending on the density of the standing of rice plants and the area of nutrition, doses and methods of fertilizer application, the optimal number of side shoots differ [1-3, 8, 9].

In this regard, the agroecosystem of rice varieties have been systematized as follows [1-3, 10]: *thinned out plantings* (sowing of 100 of viable seeds per a m²), *moderate plantings* (sowing of 300 viable seed per a m²), *highly productive crops* (sowing 500, 700 seeds per a m²) and *thickened crops* (sowing 900 germinated seeds per a m²). When adding an optimum dosage of fertilizers (N180P120 kg/ha), the largest grain yield of medium-sized rice varieties is formed in the following agroecosystem:

a) *Thinned out crops* of medium-ripening rice varieties with appearances of 6-8 lateral shoots on each rice plant; *moderate crops* with formations of 2.5-4.0 lateral shoots; *highly productive crops* - 1.8-2.5 lateral shoots; *thickened crops* with appearances of 1.1-1.6 lateral shoots on each plant;

b) for undersized samples and rice varieties, thinned out crops form the largest grain yield when there are 9-10 lateral shoots on each plant; *moderate crops* - 3.5-5.0 lateral shoots, on highly productive agroecosystem - 2-3 side shoots; *thickened crops* - with the appearance of 1.6-2.0 side shoots on each plant;

c) As for the dwarf specimen, thinned out crops produce the largest grain yield when 9-11 side shoots appear on each plant; moderate crops - 5-6 side shoots; highly productive crops - 3-4 side shoots; thickened crops - 2.1-2.5 side shoots on each rice plants.

Improving the rice varieties agroecosystem structure is important and relevant for the intensive development of rice farming [1-3]. However, the density of plant standing, formation of optimal stalk stand and its location in space, formation of plant habitus in agroecosystem and its interaction in the physiological aspect have not been sufficiently studied.

The formation of the optimal number of productive stalks in agroecosystem depends on the number of seedlings germinated and their field germination, seedling survival rate and their standing density, dose and methods of mineral fertilizers application and varietal characteristics of rice (figure; table). Thus, in medium-sized varieties and samples, the largest grain yield is formed on highly productive crops (when sowing is 500,700 viable seeds) and application N180P120 kg / ha ae. fertilizer. With an increase in the dose of fertilizer up to N240P180 kg / ha ae. grain yield has not increased, but rather decreased.

In low-growing varieties and samples of rice, the level of yield varies: in a broad-leaved variety (Lyman), the highest grain yield is formed when the optimum dosage (N180P120 kg/ha) of fertilizers is applied. On highly productive agroecosystem in varieties and samples of rice, photosynthetic activity and the

possibility of the formation of high grain yields differ. The formation of a high grain yield in varieties and samples of rice, differing in height and architectonics, is significantly affected by plant density in agrocenosis and the number of productive stems. Thus, with the introduction of the optimal dose (N180P120 kg / ha ai) of fertilizers, a high grain yield is formed under the following conditions [1-3, 11, 14, 17]:

1) A heavy yield of grain of medium-growth varieties with narrow vertically arranged leaves (Kuban 3, Krasnodar 424, Dubovsky 129) is formed in highly productive crops, with a density of rice standing on shoots of 320-400 pcs/m², before harvesting - 250-350 pcs/m², with a food area of one plant 22-30 cm², in the period of sweeping out - 25-38 cm². In such agrocenosis, the number of productive stems is 550-650 pcs / m², or 5.5-6.5 million panicles per hectare.

2) A high grain yield in medium-sized, large-leaved varieties (Marzhan, Aral 202, Togusken 1) is formed on highly productive crops, with a density of standing of 280-350 pcs / m², before harvesting 220-250 pcs/m², with 460-580 pcs./m² of productive stalks, or 4.6-5.8 million panicles per hectare.

3) Heavy yield of low varieties with large and wide leaves (Estuary) and rice samples (ECM) is ensured at a standing density of 280-380 pcs/m², before harvesting 220-310 pcs/m², with formations 550-650 pcs/m² of productive stems, or 5.5-6.5 million panicles. We called the above-mentioned crops of rice varieties, differing in height and architecture, as **high-yielding agrocenosis** [1-3, 11]. Those crops of low- and medium-growth varieties and rice samples are **representative** of specific agrocenosis. Indicators of these agrocenosis can be preliminary morphophysiological criteria (indicators) when creating models of future productive rice varieties and this should be taken into account when developing varietal technologies for their cultivation [1-3, 11].

Г. Ж. Медеуова

Қазақ мемлекеттік қыздар педагогикалық университеті, Алматы, Қазақстан

ТЫҢАЙТҚЫШТАР ДОЗАСЫНА БАЙЛАНЫСТЫ КҮРІШ СОРТТАРЫ АГРОЦЕНОЗЫНЫҢ ЖӘНЕ ДӘН ӨНІМІНІҢ ҚАЛЫПТАСУЫ

Аннотация. Биіктігі және архитектурасы бойынша айырмашылығы бар күріш үлгілері мен сорттарының жоғары дән өнімінің қалыптасуына агроценоздағы өсімдіктер тығыздылығы мен егістіктегі масақты сабақтар санының елеулі әсері бар. Атап айтқанда, оптимальды агрофон (N180P120 кг/га) жағдайында ең жоғары дән өнімі төмендегі жағдайларда қалыптасады: 1) Орта бойлы, жіңішке, тік жапырақты күріш сорттарының (Кубань 3, Краснодарский 424, Дубовский 129) ең жоғары дән өнімі себілген тұқымнан өніп шыққан өсімдіктер саны 320-400 дана/м², ору алдында 250-350 дана/м² болып, тұқымнан әдепкі өніп шыққан өскіндердің қоректену алаңы 22-30 см², масақтану кезінде 25-38 см² болғанда қалыптасады. Мұндай агроценоздағы өнімді (масақты) сабақтардың саны 550-650 дана/м², немесе гектарына 5,5-6,5 млн. масақ. 2) Орта бойлы, ірі жапырақты күріш үлгілері (4-15, КзРОС 356, Кызылординский 5) мен сорттарының (Маржан, Арал 202, Түгіскен 1) ең жоғары дән өнімі егістіктегі өсімдіктер тығыздылығы өніп шыққан кезде 280- 350 дана/м², ору алдында- 220-250 дана/м², өнімді сабақтар саны 460-580 дана/м², немесе гектарына 4,6-5,8 млн. масақты сабақтар болғанда қалыптасады. 3) Аласа бойлы, ірі жапырақты сортының (Лиман) жоғары дән өнімі егістіктегі өсімдіктер тығыздылығы өніп шыққан кезде 280-380 дана/м², ору алдында 220-310 дана/м², өнімді сабақтар саны 550-650 дана/м², немесе гектарына 5,5-6,5 млн. масақты сабақтар болғанда қалыптасады.

Түйін сөздер; күріш, сорт, агроценоз құрылымына тыңайтқыштар әсері, Тыңайт-қыштар дозасына байланысты күріш сорттары дән өнімінің құралуы.

Г. Ж. Медеуова

Казахский государственный женский педагогический университет, Алматы, Казахстан

ФОРМИРОВАНИЕ АГРОЦЕНОЗОВ СОРТОВ РИСА И УРОЖАЙНОСТЬ ЗЕРНА В ЗАВИСИМОСТИ ОТ ДОЗЫ УДОБРЕНИЯ

Аннотация. На формирование высокого урожая зерна у сортов и образцов риса, различающиеся по высоте и архитектонике, оказывают существенное влияние густота стояния растений в агроценозах, количество продуктивных стеблей. Так, при внесении оптимальной дозы (N180P120 кг/га д.в.) удобрений высокий урожай зерна формируется в следующих условиях: 1) Высокий урожай зерна среднерослых, с узким, вертикальным расположением листьев (Кубань 3, Краснодарский 424, Дубовский 129) образуется на высокопродук-

тивных посевах, при густоте стояния риса по всходам 320-400 шт/м², перед уборкой -250-350 шт/м², при площади питания одного растения 22-30 см², в период выметывания – 25-38 см². В таких агроценозах количество продуктивных стеблей равен 550-650 шт/м², или на гектаре 5,5-6,5 млн. метелок. 2) Высокий урожай зерна у среднерослых, крупнолистных сортов (Маржан, Арал 202, Тогускен 1) формируется на высокопродуктивных посевах, при густоте стояния 280-350 шт/м², перед уборкой – 220-250 шт/м², при количестве 460-580 шт/м² продуктивных стеблей, или 4,6-5,8 млн. метелок на гектаре. 3) У низкорослых, с крупными и широкими листьями сортов (Лиман) и образцов риса (ЕСМ) высокий урожай зерна формируется при густоте стояния 280-380 шт/м², перед уборкой – 220-310 шт/м², при образований 550-650 шт/м² продуктивных стеблей, или 5,5-6,5 млн. метелок.

Ключевые слова: рис, сорта, влияние удобрений на структуру агроценоза, формирование урожая сортов риса в зависимости от доз удобрений.

Information about author:

Medeuova Galiya Jumakanovna, Cand. Sc. (Agriculture), Acting Professor, Kazakh National Women Teachers Training University, Almaty, Kazakhstan; medeuova.galiya96@gmail.com; <https://orcid.org/0000-0003-3750-4758>

REFERENCES

- [1] Zhailybay K.N. Photosynthetic activity and formation of big crops of grain of grades of rice depending on agroecological factors // News of the National academy of sciences of the Republic of Kazakhstan. Series of agricultural sciences. 2017. N 4. P. 38-45. ISSN 2224-526X.
- [2] Zhailybay K.N. Agro-ecological and morpho-physiological bases of optimization of methods of application of mineral fertilizers depending on the varietal osobennosti rice // News of the National academy of sciences of the Republic of Kazakhstan. Series of agricultural sciences. 2016. N 5. P. 54-62. ISSN 2224-526X.
- [3] Zhailybay K.N., Medeuova G.Zh., Nurmash N.K. Rice Grain Quality Formation Depending on the Mineral Fertilizers Dosages // News of the National academy of sciences of the Republic of Kazakhstan. Series of biological and medical. 2018. Vol. 4, N 328. P. 103-111. ISSN 2224-5308 (Print); ISSN 2518-1629 (Online).
- [4] Fu Jianrong, Zhan Changgeng, Jinal Lina, Hasegawa M. Some issues of crops management and nutrient balance involved in high-yielding rice production // Better Crops International. 1992. Vol. 8, N 2. 8 p.
- [5] Ismunadiy M. Review of maximum yield research and high-yield technology to achieve rice self-sufficiency in Indonesia // Better Crops International. 1992. Vol. 8, N 2. 9 p.
- [6] Chae Yun Cho, Yun Jin On. Review of MYR and high-yield rice production technology in Korea // Better crops International. 1992. Vol. 8, N 2. P. 6-7.
- [7] Nguyen Vy, Pham Van Bich. Maximum yield research for maximum economic yield in Viet Nam // Better Crops International. 1992. Vol. 8, N 2. 15 p.
- [8] Urmanova Kh.U., Shchupakovskiy V.F., Nesterov A.V. Agricultural technology of growing high yields of rice // In the book: Prospects of development of rice growing in Uzbekistan. Tashkent: Publishing house of Uzbekistan, 1970. P. 50-72.
- [9] Aleshin Ye.P., Konokhova V.P. Brief reference of rice grower. M.: Agropromizdat, 1986. 253 p.
- [10] Zhailybay K.N., Sarsenbayev V.A., Ramazanova S.B., Tautenov I.A., Shermagambetov K. Agro-ecological and morpho-physiological basis of fertilizer application in rice growing in Kazakhstan (Recommendations). Almaty: Bastau, 2002. 32 p.
- [11] Zhailybay K.N. Photosynthesis and productivity of high-yielding varieties of rice (*Oryza sativa* L.) // Bulletin of agricultural science of Kazakhstan. 2006. N 1. P. 9-12.
- [12] Zhailybay K.N. Rezhim temperatury i orosheniya na zasolennykh pochvah v svyazi s sortovymi osobennostjami risa // Vestnik s.-h. nauki Kazahstana. 1996, N 7. P. 36-43.
- [13] Mustafayeva Zh.S., Kozykeyeva A.T., Kireicheva L.V., Umirzakov S.I., Zhusupova L.K. Identification of environmentally safe technology of development of solar land in production conditions of Kyzylorda region // News of the National academy of sciences of the Republic of Kazakhstan. Series of agricultural sciences. ISSN 2224-526X. Volume 6, Number 48 (2018), P.5 – 14. <https://doi.org/10.32014/2018.2224-526X.12>
- [14] Bastaubayeva Sh.O., Bekbatyrov M.B., Ospanova S.O. The biological efficiency of crop rotations and green manures to improve the productivity of irrigated lands // News of the National academy of sciences of the Republic of Kazakhstan. Series of agricultural sciences. ISSN 2224-526X. 2018. Vol. 5, N 47. P. 60-64. <https://doi.org/10.32014/2018.2224-526X.8>
- [15] Olzhabayeva A.O., Baimanov Zh.N., Daldabayeva G.T., Aldambergenova G.T., Sharmenova A.A. Improving the efficiency of using water resources on rice systems // News of the National academy of sciences of the Republic of Kazakhstan. Series of agricultural sciences. ISSN 2224-526X. 2018. Vol. 1, N 43. P. 9-21.
- [16] Ryspekov T.R., Balkozha M.A., Zhangeldina A.S. Reasons for changes of plant communities and their succession to various landscapes of Kazakhstan // News of the National academy of sciences of the Republic of Kazakhstan. Series of agricultural sciences. ISSN 2224-526X. 2018. Vol. 3, N 45. P. 54-60.
- [17] Omarchanova Zh.M., Tleuzhanova D.A., Amangeldijeva Zh.A., Baymagambetova Z.A. Analysis of development of rural territories of the Republic of Kazakhstan // News of the National academy of sciences of the Republic of Kazakhstan. Series of agricultural sciences. ISSN 2224-526X. 2018. Vol. 2, N 44. P. 37-41.

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www.nauka-nanrk.kz

ISSN 2518-1629 (Online), ISSN 2224-5308 (Print)

<http://biological-medical.kz/index.php/en/>

Редактор *М. С. Ахметова, Т. М. Апендиев, Д. С. Аленов*
Верстка на компьютере *Д. Н. Калкабековой*

Подписано в печать 07.06.2019.

Формат 60x881/8. Бумага офсетная. Печать – ризограф.
4,25 п.л. Тираж 300. Заказ 3.