

ISSN 2518-1629 (Online),
ISSN 2224-5308 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
С. Ж. Асфендияров атындағы Қазақ ұлттық медицина университеті

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

Казакский национальный медицинский
университет им. С. Д. Асфендиярова

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

Asfendiyarov
Kazakh National Medical University

**SERIES
OF BIOLOGICAL AND MEDICAL**

6 (336)

NOVEMBER – DECEMBER 2019

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Б а с р е д а к т о р

ҚР ҰҒА академигі, м. ғ. д., проф. **Ж. А. Арзықұлов**

Абжанов Архат, проф. (Бостон, АҚШ),
Абелев С.К., проф. (Мәскеу, Ресей),
Айтқожина Н.А., проф., академик (Қазақстан)
Ақшулақов С.К., проф., академик (Қазақстан)
Алшынбаев М.К., проф., академик (Қазақстан)
Бәтпенев Н.Д., проф., корр.-мүшесі (Қазақстан)
Березин В.Э., проф., корр.-мүшесі (Қазақстан)
Берсімбаев Р.И., проф., академик (Қазақстан)
Беркінбаев С.Ф., проф., (Қазақстан)
Бисенбаев А.К., проф., академик (Қазақстан)
Бишимбаева Н.Қ., проф., академик (Қазақстан)
Ботабекова Т.К., проф., корр.-мүшесі (Қазақстан)
Bosch Ernesto, prof. (Spain)
Давлетов Қ.К., ассоц.проф., жауапты хатшы
Жансүгірова Л.Б., б.ғ.к., проф. (Қазақстан)
Ellenbogen Adrian, prof. (Tel-Aviv, Israel),
Жамбакин Қ.Ж., проф., академик (Қазақстан), бас ред. орынбасары
Заядан Б.К., проф., корр.-мүшесі (Қазақстан)
Ishchenko Alexander, prof. (Villejuif, France)
Исаева Р.Б., проф., (Қазақстан)
Қайдарова Д.Р., проф., академик (Қазақстан)
Қохметова А.М., проф., корр.-мүшесі (Қазақстан)
Күзденбаева Р.С., проф., академик (Қазақстан)
Локшин В.Н., проф., корр.-мүшесі (Қазақстан)
Лось Д.А., prof. (Мәскеу, Ресей)
Lunenfeld Bruno, prof. (Израиль)
Макашев Е.К., проф., корр.-мүшесі (Қазақстан)
Миталипов Ш.М., (Америка)
Муминов Т.А., проф., академик (Қазақстан)
Огарь Н.П., проф., корр.-мүшесі (Қазақстан)
Омаров Р.Т., б.ғ.к., проф., (Қазақстан)
Продеус А.П., проф. (Ресей)
Purton Saul, prof. (London, UK)
Рахыпбеков Т.К., проф., корр.-мүшесі (Қазақстан)
Сапарбаев Мұрат, проф. (Париж, Франция)
Сарбасов Дос, проф. (Хьюстон, АҚШ)
Тұрысбеков Е.К., б.ғ.к., асс.проф. (Қазақстан)
Шарманов А.Т., проф. (АҚШ)

«ҚР ҰҒА Хабарлары. Биология және медициналық сериясы».

ISSN 2518-1629 (Online),

ISSN 2224-5308 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5546-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://biological-medical.kz/index.php/en/>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Мұратбаева көш., 75.

Г л а в н ы й р е д а к т о р

академик НАН РК, д.м.н., проф. **Ж. А. Арзыкулов**

Абжанов Архат, проф. (Бостон, США),
Абелев С.К., проф. (Москва, Россия),
Айтхожина Н.А., проф., академик (Казахстан)
Акшулаков С.К., проф., академик (Казахстан)
Алчинбаев М.К., проф., академик (Казахстан)
Батпенов Н.Д., проф. член-корр.НАН РК (Казахстан)
Березин В.Э., проф., чл.-корр. (Казахстан)
Берсимбаев Р.И., проф., академик (Казахстан)
Беркинбаев С.Ф., проф. (Казахстан)
Бисенбаев А.К., проф., академик (Казахстан)
Бишимбаева Н.К., проф., академик (Казахстан)
Ботабекова Т.К., проф., чл.-корр. (Казахстан)
Bosch Ernesto, prof. (Spain)
Давлетов К.К., ассоц. проф., ответственный секретарь
Джансугурова Л. Б., к.б.н., проф. (Казахстан)
Ellenbogen Adrian, prof. (Tel-Aviv, Israel),
Жамбакин К.Ж., проф., академик (Казахстан), зам. гл. ред.
Заядан Б.К., проф., чл.-корр. (Казахстан)
Ishchenko Alexander, prof. (Villejuif, France)
Исаева Р.Б., проф. (Казахстан)
Кайдарова Д.Р., проф., академик (Казахстан)
Кохметова А.М., проф., чл.-корр. (Казахстан)
Кузденбаева Р.С., проф., академик (Казахстан)
Локшин В.Н., проф., чл.-корр. (Казахстан)
Лось Д.А., prof. (Москва, Россия)
Lunenfeld Bruno, prof. (Израиль)
Макашев Е.К., проф., чл.-корр. (Казахстан)
Миталипов Ш.М., (Америка)
Муминов Т.А., проф., академик (Казахстан)
Огарь Н.П., проф., чл.-корр. (Казахстан)
Омаров Р.Т., к.б.н., проф. (Казахстан)
Продеус А.П., проф. (Россия)
Purton Saul, prof. (London, UK)
Рахыпбеков Т.К., проф., чл.-корр. (Казахстан)
Сапарбаев Мурат, проф. (Париж, Франция)
Сарбасов Дос, проф. (Хьюстон, США)
Турсыбеков Е. К., к.б.н., асс.проф. (Казахстан)
Шарманов А.Т., проф. (США)

«Известия НАН РК. Серия биологическая и медицинская».

ISSN 2518-1629 (Online),

ISSN 2224-5308 (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов
Министерства культуры и информации Республики Казахстан №5546-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел. 272-13-19, 272-13-18,
www.nauka-nanrk.kz / biological-medical.kz

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

Editor in chief

Zh.A. Arzykulov, academician of NAS RK, Dr. med., prof.

Abzhanov Arkhat, prof. (Boston, USA),
Abelev S.K., prof. (Moscow, Russia),
Aitkhozhina N.A., prof., academician (Kazakhstan)
Akshulakov S.K., prof., academician (Kazakhstan)
Alchinbayev M.K., prof., academician (Kazakhstan)
Batpenov N.D., prof., corr. member (Kazakhstan)
Berezin V.Ye., prof., corr. member. (Kazakhstan)
Bersimbayev R.I., prof., academician (Kazakhstan)
Berkinbaev S.F., prof. (Kazakhstan)
Bisenbayev A.K., prof., academician (Kazakhstan)
Bishimbayeva N.K., prof., academician (Kazakhstan)
Botabekova T.K., prof., corr. member. (Kazakhstan)
Bosch Ernesto, prof. (Spain)
Davletov Kairat, PhD, associate professor, executive Secretary
Dzhansugurova L.B., Cand. biol., prof. (Kazakhstan)
Ellenbogen Adrian, prof. (Tel-Aviv, Israel),
Zhambakin K.Zh., prof., academician (Kazakhstan), deputy editor-in-chief
Ischenko Alexander, prof. (Villejuif, France)
Isayeva R.B., prof. (Kazakhstan)
Kaydarova D.R., prof., academician (Kazakhstan)
Kokhmetova A., prof., corr. member (Kazakhstan)
Kuzdenbayeva R.S., prof., academician (Kazakhstan)
Lokshin V.N., prof., corr. member (Kazakhstan)
Los D.A., prof. (Moscow, Russia)
Lunefeld Bruno, prof. (Israel)
Makashev E.K., prof., corr. member (Kazakhstan)
Mitalipov Sh.M. (America)
Muminov T.A., prof., academician (Kazakhstan)
Ogar N.P., prof., corr. member (Kazakhstan)
Omarov R.T., cand. biol., prof. (Kazakhstan)
Prodeus A.P., prof. (Russia)
Purton Saul, prof. (London, UK)
Rakhypbekov T.K., prof., corr. member. (Kazakhstan)
Saparbayev Murat, prof. (Paris, France)
Sarbassov Dos, prof. (Houston, USA)
Turysbekov E.K., cand. biol., assoc. prof. (Kazakhstan)
Sharmanov A.T., prof. (USA)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of biology and medicine.

ISSN 2518-1629 (Online),

ISSN 2224-5308 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5546-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/biological-medical.kz>

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF BIOLOGICAL AND MEDICAL

ISSN 2224-5308

Volume 6, Number 336 (2019), 35 – 39

<https://doi.org/10.32014/2019.2519-1629.54>**A. A. Maui**

Kazakh National Women's Teacher Training University , Almaty, Kazakhstan.

E-mail: adilxan.maui@gmail.com**MONITORING SOY DISEASES
ON IRRIGATED LANDS OF KAZAKHSTAN**

Abstract. The flock contains data on the monitoring of soybean crops carried out by the Almaty and Zhambyl regions on the spread and development of diseases. It was found that fusarium, peronosporosis, white and gray rot are widespread soy disease, as well as ascochitosis, septariosis, anthracosis and a burn of stems. It has been shown that fusariosis occurs in all soybean crops in the Almaty region. The causative agents are fungi of the genus *Fusarium* spp. From leaf forms of diseases peronosporosis is widespread. On the Eureka-357 variety, the development of the disease was 25.7%, and the prevalence of the disease reached 100%. Infection of soybean seeds with diseases was determined. On soybean seeds, the most common pathogen is a *Fusarium* infection. It is found in all analyzed seeds of the varieties of this culture. The results of the evaluation of soybean seed dressers are presented.

Among the treated dressers, the most effective were foundazole and Maxim, where, when using these dressers, the norm was 2-3 l / t, the intensity of the development of the disease during the harvest period decreased by 7.95 and 8.2%, respectively.

Key words: soy, disinfectant, distribution, degree of disease development, germination, fusarium, white and gray rot, peronosporosis, efficiency, seeds, yield.

In recent years, in the Republic of Kazakhstan, much attention has been paid to increasing soybean production, it is being taken to increase the yield of this valuable crop, expand sown areas, and reduce crop loss from diseases. It is well known that among legumes soybean has a high content of essential nutrients. Its seeds contain up to 50% of high quality raw protein. In addition, its protein has a higher digestibility coefficient, and it is richer in valuable essential amino acids (lysine, methionine, tryptophan, etc.). By biological value, soy is compared with the protein of meat, milk, eggs, the fat content of 25% and no worse than butter from cow's milk. Soy also contains minerals vitamins [1]. All this allows you to use it for food, technical and feed purposes [2].

Soybean is also of great importance in increasing soil fertility. Due to nodule bacteria, the arable layer is enriched with organic nitrogen. In addition, it is a good precursor for many [3,4] crops.

Being a culture of multifaceted use, soy is unparalleled in its universality of application. Already now it is used for the manufacture of more than 400 types of industrial products.

As you can see, the economic value of soybeans is multifaceted, therefore, a comprehensive increase in the size of sown areas in southeast Kazakhstan and its advancement to the east (East Kazakhstan region), west (Aktyubinsk, West Kazakhstan region) and north (Kostanay region) countries is an important step in the intensification of agricultural production. The introduction of this crop in the irrigated lands of the south, southeast, east, west and north of the republic that meets the biological and biological requirements of soy according to the soil and climatic conditions will make it possible to increase the productivity of each irrigated hectare, the payback of costs and the productivity of animal husbandry.

Currently, sown area under soybean reaches 150 thousand ha, which is not enough for Kazakhstan. The reserves for obtaining raw materials for the production of soy protein in the republic are essentially unlimited. Kazakhstan has every opportunity to bring clean soybean crops to 1.0 million ha.

In the republic, in leading equitable farms, subject to all the rules of agricultural technology, they get a high yield. So, in 2000 in the PC named after D.A. Kunaev, Talgar district, Almaty region, the soybean grain yield was 46.5c / ha, in the same year, the Aktyubinsk regional station received a crop of 27.09 / ha, and in 2015 on an experimental base of the Kazakh Research Institute of Agriculture and Plant Growing, the yield per hectare reached 67 , 5ts / ha. At the same time, in recent years, there has been a tendency to an increase in yield losses of this valuable crop due to the spread of various harmful organisms in its crops, which include infectious diseases that affect soybeans during different periods of vegetation.

Currently, according to scientists around the world, more than three hundred diseases have been discovered on soy. For example, in the USA, 25 pathogens are a constant threat to this culture, of which 19 are fungal, 3 are bacterial and 3 are viral. In China, out of the 8 most common diseases, fungal infections are 6. In Russia, only 32 soy pathogens are known, in Ukraine - 23. In the Russian Federation, among the 6-7 main soy pathogens, as a rule, 1-2 bacterial [5] are constantly called.

Soy diseases annually cause enormous damage to the crop of this valuable crop. So, in years with normal weather conditions in the Far East [6,7], due to damage to plants by various diseases, seed yields decrease by 20-30%, and in years with heavy rainfall by 50% or more.

We are in 2013-2017. Together with the staff of the leguminous crops department of the Kazakh Research Institute of Agriculture and Plant Growing, the phytosanitary condition of soybean crops was studied. The spread of diseases and the species composition of pathogens was studied by route surveys of soybean crops in Almaty and Zhambyl regions, and seed contamination was determined according to generally accepted methods [8, 9].

More than 20 diseases were registered on soybean plants in the republic: 15 fungal, 3 viral and 3 bacterial diseases. Among soy diseases in the republic, the most common and harmful are fusarium, peronosporosis, white and gray rot, and also in some regions of the country - phomopsis.

Fusarium is especially widespread in recent years. The causative agents are fungi of the genus *Fusarium* spp. The fungus infects seedlings and causes plants to wilt. So, the death of shoots from *Fusarium* infections in a cold and protracted spring can reach 30-50%. They were most affected by soybean in those farms where crops were thickened, clogged and crop rotation was not observed. So, in certain fields of the farm "Amangeldi" of the Koksuy region, the damage to plants by the disease reached 50% [11]. In other farms on soybean crops, foci were found where 15-25% of plants were affected. In this regard, some fields had to be reseeded.

In wet years, white and gray rot spread strongly on soybeans (they were observed everywhere in 2013, 2016 and 2017). Of the diseases affecting mainly plant leaves, downy mildew is widespread. The degree of damage in wet years reaches up to 100%, which leads to a decrease in the assimilation surface of the leaves, as a result of this pathogen can reduce the yield of this culture to 60-70%.

Surveys have shown that in a number of farms the development of epiphytoses of white, gray rot and peronosporosis or downy mildew was promoted not only by weather conditions, but also by violations of the technology of cultivation of the crop. Some farms do not comply with crop rotation and return soybeans to their former place in two to three years or even earlier. There is a farm culture cultivated in monoculture. This leads to massive accumulation of fungal infection in the soil.

Farm "Turgen" of Enbekshikazakhsky district soybean crops were heavily clogged with sow thistle and annual weeds, where 70% of the field was affected by plants affected by white and gray rot. Ayyr-Shir LLP of Talgar region, where soya is paid enough attention, is grown on a high agricultural background, there were no diseases on the crop during the growing season or they appeared to a weak degree before harvesting.

Zoned varieties of local and foreign selection, as well as varieties and varietal samples that are in a competitive test, are not resistant to diseases affecting soybeans (table 1).

Seeds serve as a source of infection of plants with dangerous infectious diseases. They can preserve the causative agents of many infectious diseases, be a source of renewal for the next year; with seeds, pathogens can be transferred to new areas where they did not exist before, i.e. are important in the migration of pathogenic microorganisms and the spread of infectious diseases of soy. Seed infection in soybeans is very harmful, its development on seedlings leads to the death of plants. Plants affected by *Fusarium* root rot lag behind in growth, do not form at all or form small beans that produce unobtrusive seeds. As a result of monitoring soybean crops, we selected 17 batches of soybean seeds for phyto-examination in 8 farms of the Almaty region. Its results are presented in table 2.

Table 1 – Infection of commercial soybean varieties with diseases during the growing season of Kazakh scientific research institute of protection and crop production 2013-2018

Class	Origin of class	Amazed, %		Growth, %
		fusarium	sclerotiniosis	peronosporosis
Kazakhstan 200	Kazakhstan	6,5	5,5	13,3
Almaty	Kazakhstan	3,5	0,0	9,5
Savage	France	1,0	0,0	3,0
Sting	Kazakhstan	12,5	4,5	13,5
Tazhin	Netherlands	2,5	0,0	21,1
Eureka-357	Moldova	7,5	4,0	25,7
Zen	China	9,5	0,0	17,0
SibNISH-1	Russia	27,0	10,0	13,5
Swallow	Kazakhstan	7,5	1,5	8,0
Zhansaya	Kazakhstan	11,5	6,5	23,0
Vita	Kazakhstan	8,5	7,0	17,8

Table 2 – Infection of soybean seeds with diseases in the conditions of Almaty region

Disease	Weighted average percentage of infection, %		
	2015	2016	2017
Fusarium	0,5	1,0	7,5
Peronosporosis	0,7	1,2	13,0
Septariosis	2,0	0,2	3,5
Burnstems	–	0,1	0,5
Ascochitosis	1,5	1,1	6,0
Alternariosis	0,1	0,5	5,5

We must not underestimate the danger of penetration into the republic of pathogens that are absent in our country with imported foreign seeds. A similar thing already happened when phomopsis entered Kazakhstan from abroad with infected seed material.

In addition, a number of diseases are described in different countries of the world that we have not yet discovered, however, judging by the biology of the causative agents of the diseases and given the similar climatic conditions, there is reason to believe that these diseases may eventually appear on our soy plantations.

By us during 2017-2018. At the experimental base of the Kazakh Scientific Research Institute of Agriculture and Plant Growing, as well as the laboratory of the Kazakh National Women's Pedagogical University, the influence of protectants on seed germination and disease emergence was studied.

To improve seed health from fusarium, sclerotiniosis, ascochitosis, peronosporosis and other diseases, we studied the following drugs: foundationazole (3l / t), rovril (3kg / t), apron (6kg / t), derazole (3kg / t), tachigaren (6kg / t), maxim (2l / t), scarlet (2l / t), vitalon (3l / t). Control seeds were not etched. TMTD-4kg / t was used as a reference. Etching was carried out one to two days before sowing. Variety Zhapak-Saz. The area of each plot was 25 sq.m., the repetition of 4-fold. When studying the effect of dressing agents on seed germination, the latter were sown in laboratory and field conditions, all preparations had a positive effect on seed germination (table 3), the highest results were noted in variants with foundationazole, tachigaren, maxim, scarlet and vitalon. Their laboratory germination rate was 85.9, respectively; 85.5; 85.0; 84.5; 83.0% and exceeded control by 8, respectively; 7.5; 7; 5.5 and 5%, the variant with ethanol is 4; 3.5; 3.0; 1.5 and 1.0%. The preparations acted favorably on the increase in field germination of seeds in comparison with the control by 1.3–13.8%.

Table 3 – Efficiency of soybean seed dressers (grade Zhalspak-Saz, 2017-2018)

Variant	Consumption rate, kg, l / t	Laboratory germination, %	Field germination, %	The defeat of Fusarium rot,%				Harvest, t / ha
				seedlingphase		beforecleaning		
				affected plants	disease-progression	affected plants	disease-progression	
Control without treatment	–	78,0	59,2	32,0	11,4	58,5	19,2	30,5
TMTD (standard)	4,0	82,0	63,7	22,1	5,04	42,0	15,2	33,2
Tachigaren	6,0	83,5	72,0	6,4	0,6	25,0	2,5	35,5
Apron	6,0	79,5	61,5	23,4	7,6	56,5	17,5	32,8
Fundazole	3,0	83,0	66,0	8,1	0,3	38,5	11,25	33,7
Scarlet	2,0	83,0	66,7	16,4	1,4	52,5	17,3	33,0
Derazole	3,0	82,0	63,7	20,3	5,3	40,0	17,8	33,0
Rovral	3,02,0	85,0	66,5	9,6	1,0	35,0	11,0	34,3
Maksim	2,0	82,5	65,5	9,1	0,5	35,5	11,0	34,0
Vitalon	2,0	84,5	67,0	10,0	0,7	39,0	12,0	33,9

On average, over two years in the variants with the use of foundationazole (3 l / t), maxim (2 l / t) and vitalon (3 l / t), the development of fusarium rot was 0.3; 0.5 and 0.7%, which is less than the control by 11.1; 10.9 and 10.7% and was lower than in versions with a standard (TMTD, 4kg / t) by 5.4; 4.9 and 4.7%. Positive results were obtained when treating soybean seeds with a scarlet (2 l / t) and tachigaren (6 kg / t). A similar pattern was observed during the harvest period. So, during the pre-sowing treatment of soybean seeds with rovrал in the norm of 3 kg per 1 ton of seeds, the degree of damage to plants with fusarium rot decreased by 8.12%. In the fight against fusarium rot, a high effect was exerted by derosal and foundationazole at a rate of 3 l / t, the intensity of the development of the disease during the harvest period decreased by 2.4 and 7.97%, respectively.

The use of dressing agents has increased soybean productivity. So, the yield in variants with the use of foundationazole and Maxim, respectively, amounted to 33.7; and 34.0 t / ha, which is 3.2 and 0.8 t / ha more than in the control (30.5 t / ha) and 0.7 and 0.8 t / ha higher compared to seed treatment TMTD (4 kg / t).

Thus, among the studied protectants, the most effective were fundazole and Maxim. The use of these drugs helps to increase field germination by 6.3 and 6.8%, reduce the prevalence and development of fusarium rot in the seedling phase by 22.9-23.9% and 2.4-17.7%.

Ә. Ә. Мәуи

Қазақ Ұлттық қыздар педагогикалық университеті, Алматы, Қазақстан

КАЗАКСТАННЫҢ СУАРМАЛЫ ЕГІС АЛҚАПТАРЫНДАҒЫ ҚЫТАЙ БҰРШАҒЫНЫҢ АУРАЛАРЫНА МОНИТОРИНГ

Аннотация. Қазақстанның қытай бұршағы егілетін суармалы егіс алқаптарына жүргізілген мониторинг нәтижесінде осы дақылдың ауруларының таралуы мен дамуы анықталды. Кең тараған аурулардың түрлеріне мыналар жағада, фузариоз, ақ және сұр шіріктері және жапырақтарында көптеген дақ аурулары кездеседі. Алматы облысының егіс алқаптарында өте қауіпті және зиянды фузариоз ауруы кең тараған, ал жапырақтарында пероноспороз патогені кең көлемде кездеседі. Эврика-357 сортында аталған аурудың даму деңгейі 25,7% жеткен. Сонымен бірге қытай бұршағының тұқыммен таралатын ауру қоздырғыштары анықталды. Қытай бұршағы тұқымымен берілетін ауру қоздырғыштарын препараттармен дәрілеудің тиімділігі келтірілген. Максим және фундазол препараттарының тұқым ауруына тиімділігі жоғары екені көрсетілген.

Түйін сөздер: Қытай бұршағы, патоген, таралуы, дамуы, аурулар, өнгіштігі, фузариоз, ақ және сұр шіріктері, пероноспороз, тиімділік, тұқым, өнім.

А. А. Мауи

Казахский национальный женский педагогический университет, Алматы, Казахстан

МОНИТОРИНГ БОЛЕЗНЕЙ СОИ НА ОРОШАЕМЫХ ЗЕМЛЯХ КАЗАХСТАНА

Аннотация. В статье приводятся данные по мониторингу посевов сои, проведенного Алматинской и Жамбылской областях по распространению и развитию болезней. Установлено, что широко распространенной болезнью сои являются фузариоз, пероноспороз, белая и серая гнили, а также встречаются аскохитоз, септариоз, антрокноз и ожог стеблей. Показано, что на всех посевах сои в Алматинской области встречается фузариоз. Возбудители грибы рода *Fusarium* spp. Из листовых форм болезней широко распространен пероноспороз. На сорте Эврика-357 развитие болезни составило 25,7%, а распространенность болезни достигало 100%. Определена зараженность семян сои болезнями. На семнах сои наиболее распространенным возбудителем является фузариозная инфекция. Она встречается во всех анализированных семенах сортов этой культуры. Представлены результаты оценки протравителей семян сои.

Среди изученных протравителей наиболее эффективными оказались фундазол и Максим, где при применении указанных протравителей в норме 2-3л/т, интенсивность развития болезни в период уборки уменьшилась соответственно на 7,95 и 8,2%.

Ключевые слова: соя, протравитель, распространение, степень развития болезни, всхожесть, фузариоз, белая и серая гнили, пероноспороз, эффективность, семена, урожай.

Information about author:

Maui A. A., DSc., professor, Kazakh National Women's Teacher Training University, Almaty, Kazakhstan; adilxan.maui@gmail.com; <https://orcid.org/0000-0002-4518-8743>

REFERENCES

- [1] Balakai, G.T. Soya: agricultural technology, processing. Rostov on the Don. Phoenix, 2003. -- 160 p.
- [2] Karyagin Yu.G. Recommendations on soybean cultivation on irrigated lands of the south and southeast of Kazakhstan. Alma-Ata 1974. P. 26
- [3] V.P. Blackhead. Legumes in Kazakhstan. // Alma - Ata: Kaynar ", 1974. - S.47 - 51.
- [4] E.V. Bakaev, V.P. Blackhead. Soya in Kazakhstan. Alma-Ata. Publishing house "Kazselkhozgiz". 1963.p.33.
- [5] A.M. Ovchinnikov. Mushroom soy diseases // Diseases and pests of soy in the south of the Far East and measures to combat them. Vladivostok, 1971. - P. 5 -72.
- [6] I.N. Abramov. Fungal diseases of soybeans in the Far East. Vladivostok. 1931. p. 84.
- [7] L.K. Dubovitskaya. Root rot of soybeans in the Amur region and the rationale for measures to combat it. Thesis. – Blagoveshchensk. 1980.138 p.
- [8] GOST 12038-84. Seeds of crops. Germination determination methods. –М. : Publishing house of standards. 1991.57 p.
- [9] GOST 12044-93. Seeds of crops. Methods for the determination of disease infection. - М. : Publishing house of standards. 2012.58 p.
- [10] A.A. Abubakirova, A.D. Daulbai, A.A. Ospanova, R.A. Abildaeva, S.Zh. Lesbekova. "Studies of the biological characteristics of the spread of pathogenic fungi in soybean plants." Proceedings of the National Academy of Sciences of the Republic of Kazakhstan. A series of biological and medical. No. 2. 2017. S. 88-93.
- [11] A.Sh. Raushanova, R.A. Iskendiroya. "Mushroom diseases of soy in the conditions of southeastern Kazakhstan." Proceedings of the National Academy of Sciences of the Republic of Kazakhstan. A series of biological and medical. No. 2. 2018. S. 10-14.

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

ISSN 2518-1629 (Online), ISSN 2224-5308 (Print)

<http://biological-medical.kz/index.php/en/>

Редактор *М. С. Ахметова, Т. М. Апендиев, Д. С. Аленов*
Верстка на компьютере *Д. Н. Калкабековой*

Подписано в печать 06.12.2019.

Формат 60x881/8. Бумага офсетная. Печать – ризограф.
2,75 п.л. Тираж 300. Заказ 6.