

ISSN 2518-1629 (Online),
ISSN 2224-5308 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
С. Ж. Асфендияров атындағы Қазақ ұлттық медицина университеті

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Қазақстан Республикасының Ғылым Академиясының
С. Ж. Асфендияров атындағы Қазақ ұлттық медицина университеті

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Asfendiyarov
Kazakh National Medical University

**SERIES
OF BIOLOGICAL AND MEDICAL**

2 (338)

MARCH – APRIL 2020

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Б а с р е д а к т о р

ҚР ҰҒА академигі, м. ғ. д., проф.

Ж. А. Арзықұлов

Абжанов Архат, проф. (Бостон, АҚШ),
Абелев С.К., проф. (Мәскеу, Ресей),
Айтқожина Н.А., проф., академик (Қазақстан)
Акшулаков С.К., проф., академик (Қазақстан)
Алшынбаев М.К., проф., академик (Қазақстан)
Бәтпенев Н.Д., проф., корр.-мүшесі (Қазақстан)
Березин В.Э., проф., корр.-мүшесі (Қазақстан)
Берсімбаев Р.И., проф., академик (Қазақстан)
Беркінбаев С.Ф., проф., (Қазақстан)
Бисенбаев А.К., проф., академик (Қазақстан)
Бишимбаева Н.Қ., проф., академик (Қазақстан)
Ботабекова Т.К., проф., корр.-мүшесі (Қазақстан)
Bosch Ernesto, prof. (Spain)
Давлетов Қ.К., ассоц.проф., жауапты хатшы
Жансүгірова Л.Б., б.ғ.к., проф. (Қазақстан)
Ellenbogen Adrian, prof. (Tel-Aviv, Israel),
Жамбакин Қ.Ж., проф., академик (Қазақстан), бас ред. орынбасары
Заядан Б.К., проф., корр.-мүшесі (Қазақстан)
Ishchenko Alexander, prof. (Villejuif, France)
Исаева Р.Б., проф., (Қазақстан)
Кайдарова Д.Р., проф., академик (Қазақстан)
Кохметова А.М., проф., корр.-мүшесі (Қазақстан)
Күзденбаева Р.С., проф., академик (Қазақстан)
Локшин В.Н., проф., корр.-мүшесі (Қазақстан)
Лось Д.А., prof. (Мәскеу, Ресей)
Lunenfeld Bruno, prof. (Израиль)
Макашев Е.К., проф., корр.-мүшесі (Қазақстан)
Миталипов Ш.М., (Америка)
Муминов Т.А., проф., академик (Қазақстан)
Огарь Н.П., проф., корр.-мүшесі (Қазақстан)
Омаров Р.Т., б.ғ.к., проф., (Қазақстан)
Продеус А.П., проф. (Ресей)
Purton Saul, prof. (London, UK)
Рахыпбеков Т.К., проф., корр.-мүшесі (Қазақстан)
Сапарбаев Мұрат, проф. (Париж, Франция)
Сарбасов Дос, проф. (Хьюстон, АҚШ)
Тұрысбеков Е.К., б.ғ.к., асс.проф. (Қазақстан)
Шарманов А.Т., проф. (АҚШ)

«ҚР ҰҒА Хабарлары. Биология және медициналық сериясы».

ISSN 2518-1629 (Online), ISSN 2224-5308 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.).

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5546-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28; 219, 220 бөл.; тел.: 272-13-19, 272-13-18;
<http://biological-medical.kz/index.php/en/>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2020

Типографияның мекенжайы: «NurNaz GRACE», Алматы қ., Рысқұлов көш., 103.

Г л а в н ы й р е д а к т о р
академик НАН РК, д.м.н., проф.
Ж. А. Арзыкулов

Абжанов Архат, проф. (Бостон, США),
Абелев С.К., проф. (Москва, Россия),
Айтхожина Н.А., проф., академик (Казахстан)
Акшулаков С.К., проф., академик (Казахстан)
Алчинбаев М.К., проф., академик (Казахстан)
Батпенов Н.Д., проф., чл.-корр. (Казахстан)
Березин В.Э., проф., чл.-корр. (Казахстан)
Берсимбаев Р.И., проф., академик (Казахстан)
Беркинбаев С.Ф., проф. (Казахстан)
Бисенбаев А.К., проф., академик (Казахстан)
Бишимбаева Н.К., проф., академик (Казахстан)
Ботабекова Т.К., проф., чл.-корр. (Казахстан)
Bosch Ernesto, prof. (Spain)
Давлетов К.К., ассоц. проф., ответственный секретарь
Джансугурова Л. Б., к.б.н., проф. (Казахстан)
Ellenbogen Adrian, prof. (Tel-Aviv, Israel),
Жамбакин К.Ж., проф., академик (Казахстан), зам. гл. ред.
Заядан Б.К., проф., чл.-корр. (Казахстан)
Ishchenko Alexander, prof. (Villejuif, France)
Исаева Р.Б., проф. (Казахстан)
Кайдарова Д.Р., проф., академик (Казахстан)
Кохметова А.М., проф., чл.-корр. (Казахстан)
Кузденбаева Р.С., проф., академик (Казахстан)
Локшин В.Н., проф., чл.-корр. (Казахстан)
Лось Д.А., prof. (Москва, Россия)
Lunenfeld Bruno, prof. (Израиль)
Макашев Е.К., проф., чл.-корр. (Казахстан)
Миталипов Ш.М., (Америка)
Муминов Т.А., проф., академик (Казахстан)
Огарь Н.П., проф., чл.-корр. (Казахстан)
Омаров Р.Т., к.б.н., проф. (Казахстан)
Продеус А.П., проф. (Россия)
Purton Saul, prof. (London, UK)
Рахыпбеков Т.К., проф., чл.-корр. (Казахстан)
Сапарбаев Мурат, проф. (Париж, Франция)
Сарбасов Дос, проф. (Хьюстон, США)
Турысбеков Е. К., к.б.н., асс. проф. (Казахстан)
Шарманов А.Т., проф. (США)

«Известия НАН РК. Серия биологическая и медицинская».

ISSN 2518-1629 (Online), ISSN 2224-5308 (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов
Министерства культуры и информации Республики Казахстан №5546-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28; ком. 219, 220; тел. 272-13-19, 272-13-18;

<http://biological-medical.kz/index.php/en/>

© Национальная академия наук Республики Казахстан, 2020

Адрес типографии: «NurNazGRACE», г. Алматы, ул. Рыскулова, 103.

Editor in chief

Zh.A. Arzykulov,
academician of NAS RK, Dr. med., prof.

Abzhanov Arkhat, prof. (Boston, USA),
Abelev S.K., prof. (Moscow, Russia),
Aitkhozhina N.A., prof., academician (Kazakhstan)
Akshulakov S.K., prof., academician (Kazakhstan)
Alchinbayev M.K., prof., academician (Kazakhstan)
Batpenov N.D., prof., corr. member (Kazakhstan)
Berezin V.Ye., prof., corr. member (Kazakhstan)
Bersimbayev R.I., prof., academician (Kazakhstan)
Berkinbaev S.F., prof. (Kazakhstan)
Bisenbayev A.K., prof., academician (Kazakhstan)
Bishimbayeva N.K., prof., academician (Kazakhstan)
Botabekova T.K., prof., corr. member (Kazakhstan)
Bosch Ernesto, prof. (Spain)
Davletov Kairat, PhD, associate professor, executive Secretary
Dzhansugurova L.B., Cand. biol., prof. (Kazakhstan)
Ellenbogen Adrian, prof. (Tel-Aviv, Israel),
Zhambakin K.Zh., prof., academician (Kazakhstan), deputy editor-in-chief
Ishchenko Alexander, prof. (Villejuif, France)
Isayeva R.B., prof. (Kazakhstan)
Kaydarova D.R., prof., academician (Kazakhstan)
Kokhmetova A., prof., corr. member (Kazakhstan)
Kuzdenbayeva R.S., prof., academician (Kazakhstan)
Lokshin V.N., prof., corr. member (Kazakhstan)
Los D.A., prof. (Moscow, Russia)
Lunefeld Bruno, prof. (Israel)
Makashev E.K., prof., corr. member (Kazakhstan)
Mitalipov Sh.M. (America)
Muminov T.A., prof., academician (Kazakhstan)
Ogar N.P., prof., corr. member (Kazakhstan)
Omarov R.T., cand. biol., prof. (Kazakhstan)
Prodeus A.P., prof. (Russia)
Purton Saul, prof. (London, UK)
Rakhypbekov T.K., prof., corr. member (Kazakhstan)
Saparbayev Murat, prof. (Paris, France)
Sarbasov Dos, prof. (Houston, USA)
Turysbekov E.K., cand. biol., assoc. prof. (Kazakhstan)
Sharmanov A.T., prof. (USA)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of biology and medicine.
ISSN 2518-1629 (Online), ISSN 2224-5308 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty).

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5546-Ж, issued 01.06.2006.

Periodicity: 6 times a year.

Circulation: 300 copies.

Editorial address: 28, Shevchenko str. of. 219, 220, Almaty, 050010; tel. 272-13-19, 272-13-18;

<http://biological-medical.kz/index.php/en/>

© National Academy of Sciences of the Republic of Kazakhstan, 2020

Address of printing house: «NurNaz GRACE», 103, Ryskulov str, Almaty.

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF BIOLOGICAL AND MEDICAL

ISSN 2224-5308

Volume 1, Number 337 (2020), 12 – 19

<https://doi.org/10.32014/2020.2519-1629.8>

UDC 578.832.1:578.4

IRSTI 34.25.39; 34.25.21

K. Kh. Zhumatov, A. I. Kydyrmanov, M. Kh. Sayatov

Scientific and Production Center of Microbiology and Virology, Almaty, Kazakhstan.

E-mail: kainar60@yahoo.com, kydyrmanov@yandex.kz, sayatov37@mail.ru

INFLUENZA D VIRUSES - PATHOGENS FORMING A NEW GENUS IN THE ORTHOMYXOVIRIDAE FAMILY

Abstract. Influenza pathogens belong to the Orthomyxoviridae family and are divided into genera: Influenzavirus A, B, C, D, as well as Quaranjavirus, Thogotovirus, and Isavirus. For the first time, the influenza D virus was isolated from swine nasal swabs in 2011 in the United States, and its widespread distribution among cattle in France, China, Italy, Ireland, Japan, and several African countries, as well as its ability to infect ferrets, guinea pigs, is further shown. Antibodies to influenza D virus are found in the blood serum of horses, sheep, goats, and in people who have been in contact with cattle. The RNA genome of the influenza D virus is represented by seven fragments responsible for the synthesis of nine proteins. The longest three segments encode for polymerases PB2, PB1, and P3; the fourth and fifth segments encode for hemagglutinin-esterase fusion protein – HEF and nucleoprotein – NP, respectively. The sixth fragment is involved in the synthesis of membrane polypeptides DM1 and DM2, which, in accordance, lines the viral membrane from the inside and performs the function of proton channels. The seventh segment encodes the non-structural protein NS1 and the nuclear export protein NEP; NS1 helps to neutralize cellular interferon and NEP mediates the nuclear export of ribonucleoprotein. Three phylogenetic lines of the influenza virus D are described – D/OK, D/660, and D/Japan, which must be taken into account when preparing vaccines. It is concluded that from its epidemiological, pathological and biological characteristics, the potential ability to cause disease in humans and be transmitted from person to person, new, more in-depth studies are required using ecological-virological and molecular genetic methods.

Key words: virus, influenza D, genome, variability, cattle, phylogenesis, clade, HEF fusion protein, serology.

Characteristics of the Orthomyxoviridae family. Influenza occupies one of the first places among infectious diseases in terms of the number of species involved in the infectious process and is characterized by global distribution and high economic and social significance [1].

All currently known influenza pathogens belong to the family *Orthomyxoviridae* and are divided into genera: *Influenzavirus A, B, C, D*, as well as *Quaranjavirus, Thogotovirus* and *Isavirus*; the last two infect rabbit-like mammals and salmon fish [2,3,4]. Representatives of *Quaranjavirus* found among both invertebrates (ticks) and vertebrates (water birds) hosts [5].

Influenza A viruses are widespread in the environment and infect humans, mammals and birds. The global spread of influenza A pathogens is due to unique variability, which is based on point mutations and recombinations of eight segments of the genome. The classification of influenza A viruses is determined by a combination of the known subtypes of surface antigens of hemagglutinin (HA) and neuraminidase

(NA) – H1N1, H3N2, H5N1, H7N7, etc. [6]. In Kazakhstan, as a result of long-term studies of influenza A virus ecology, eight different subtypes of this pathogen were identified [7,8,9,10,11,12,13].

The causative agents of influenza A along with influenza B viruses, cause annual epidemics, accompanied by 3-5 million cases of severe morbidity and about 300,000 deaths, with a mortality rate reaching 16 % and 10 %, respectively [14,15].

Modern epidemic influenza B viruses are phylogenetically and antigenically divided into two lines - Yamagata-like and Victoria-like, which are easily differentiable in the hemagglutination inhibition assay (HI) [16]. The natural host of influenza B virus is human but sporadic infections of pheasants, horses, and dogs have been reported in 1960-1980 [17,18,19]. Further, this virus was isolated from a seal (*Phoca vitulina*) in the Netherlands in 1999, the B/Seal/Netherlands/1/99 [20]. Antibodies to influenza B virus were detected in pig sera in China in 2015, their sensitivity to experimental infection was also shown [21].

Representatives of the genus C predominantly infect the children's contingent and differ significantly in structural characteristics from A and B viruses. They possess an RNA genome consisting of seven fragments, one of which encodes hemagglutinin-esterase (HEF) synthesis, which combines the functions of HA and NA in the virion. Another feature of the influenza C virus (ICV) is the ability to infect pigs, which was first demonstrated in China in 1982 [22], in addition to this, antibodies to it were detected in 2015 in the horse sera of in the USA [23].

The Orthomyxoviridae family is represented by the so-called newly emerging infectious diseases, the number of which is growing in the world, and many of them pose a serious threat to wildlife, domestic animals, and public health [24,25].

In April 2011, B.M. Hause et al. [26] in Oklahoma (USA), during the virological study of nasal swabs from pigs, isolated an influenza virus with an RNA genome consisting of 7 fragments that are 50% structurally similar in amino acid sequence to ICV. Originally designated as C/Oklahoma/1334/2011, it diverged from ICV in phylogenetic analysis to the same extent as influenza A viruses differ from representatives of genus B. No cross-reactivity was observed between the new isolate and ICV in HI. Antibodies to this strain were found in 9.5 % and 1.3 % of pig sera and staff, respectively; in addition, it was transmitted to healthy animals through direct contact and was able to infect pigs, ferrets, and guinea pigs in the experiment. Cell tropism of the new virus was superior to that of ICV, while conservative enzymatic and divergent receptor-binding sites were detected in the HEF fusion protein. In experiments with two human ICVs and two newly isolated swine and cattle viruses, no reassortment and production of viable generation were found. In experiments with specific polyclonal antibodies, cross-recognition was not observed in agar gel immunodiffusion. Based on the data obtained, the isolated virus was assigned to the new genus D [27], which was adopted by the decision of the International Committee on Taxonomy of Viruses (ICTV) in 2018 [28]. Subsequent molecular genetic studies have identified several characteristics of the causative agent of influenza D.

Structural features of the influenza D virus. The RNA genome of the influenza D virus (IDV) is represented by seven fragments responsible for the synthesis of 9 proteins. The longest three segments encode polymerases PB2, PB1, and P3, the fourth – protein HEF, the fifth – NP. The sixth fragment is involved in the synthesis of membrane polypeptides DM1 and DM2, one of which underlines the viral membrane from the inside, the other performs the function of proton channels [29]. The seventh segment encodes the non-structural protein NS1, and the nuclear export protein NEP; NS1 helps neutralize cellular interferon, NEP mediates the nuclear export of ribonucleoprotein [30]. IDV, like ICV, uses cellular 9-O-acetylated sialic acid as its receptor but exhibits a broader cellular and host tropism [31]. The reason for this lies in the structural features of the molecule of the HEF monomer, consisting of two subunits HEF1 and HEF2. As revealed by comparative X-ray crystallographic study and following the domain nomenclature previously used for ICV, H. Song et al. [31] divided the structure of IDV HEF into three domains: a receptor-binding domain (R), an esterase domain (consisting of subdomains E1, E' and E2) and a fusion domain (consisting of subdomains F1, F2, and F3). The general structure and structural folds of the individual HEF subdomains of both viruses were similar. E domains are the most conserved; the F1 and F2 subdomains are less identical, while the F3 subdomain contains the fusion peptide, which is essential for the viral membrane fusion.

The E domain of IDV HEF, harboring the receptor-destroying enzyme (RDE) activity, has a hydrolase fold that is highly similar to that of ICV HEF. The active-site architecture of the HEF sialate-9-O-acetylerase is fully conserved in ICV and IDV HEF and has oxyanion hole. The pocket is

extremely conserved not only in IDV HEF and ICV HEF but also in some nidovirus hemagglutinin-esterase (HE) proteins such as bovine coronavirus, porcine torovirus and bovine torovirus HE [32]. Phylogenetic analysis also shows the relationship between HEF and HE to be much closer than the relationship between HEF and HA or hemagglutinin-neuraminidase (HN) proteins, implying common ancestral origins.

Epidemiology of influenza D. Following the first report on the isolation of IDV in 2013, L. Ferguson et al. [33] in 2014 recovered 15 IDVs from surveillance of bovine herds in Mississippi belonging to two genetic lines. Based on the serological analysis of “archived” sera, it was shown that the pathogen has been circulating in cattle stock in this region since at least 2014. Active transmission of the virus has been detected in places where newborn, weaned, and comingled calves were maintained. The detection rate of IDV in RT-PCR was 2.4 % of 82 healthy calves, which was significantly higher than in Chinese Shandong – 0.7 % of 453 calves [34], which indicates its possible participation in the etiology of respiratory cattle disease, as an infection with many potential pathogens. Continuing his research, L. Ferguson et al. [29] conducted an experimental intranasal infection of three dairy calves with the influenza virus D/bovine/Mississippi/C00046N/2014 followed by replanting of a seronegative contact calf. Seroconversion and the presence of the virus in the respiratory tract were detected in all animals. The infection of ferrets with nasal discharge from calves has not yielded results. These data are consistent with Koch's postulates and confirm the hypothesis that cattle are a natural reservoir of the virus.


In subsequent years, IDVs were found in cattle in France, China, Italy, and Japan [35,34,36,37]. E. Collin et al. [38] studied 208 biological samples from cattle from 12 US states in real-time PCR and found 10 positives for IDV samples, using passages on cell cultures, they isolated six IDV strains.

O. Flynn et al. [38] examined 320 “archival” samples in the form of nasal swabs taken from cattle in Ireland during 2014–2016. Using RT-PCR on IDV and Sanger sequencing, five partial RNA sequences suitable for phylogenetic analysis were obtained from 18 positive samples.

In Japan, according to a national serological study, the average level of seropositivity in cattle for the influenza virus is 30%, and the values vary depending on the region [37,40]. At the same time, a molecular epidemiological analysis of cattle samples with symptoms of respiratory disease collected from 23 different farms in this country revealed 2.1 % (8/377) influenza D infection [41].

E. Salem et al. [42] conducted a serological search for antibodies to influenza C and D viruses among ruminants and camelids in Africa in 1991–2015, and for this purpose, they collected 2083 blood serum samples from a cattle, pigs, small ruminants, and dromedary camels in Morocco (n=200), Togo (n=540), Côte d'Ivoire (n=203), Benin (n=308) and Kenya (n=1231). The percentage of positive samples from cattle in Morocco was 35 %, Togo – 10.4 %, Côte d'Ivoire – 0 %, Benin – 1.9 %, Kenya – 0 %. Noteworthy is the high percentage (99 %) of positive blood serum from camels in Kenya (287 out of 293), but the serum was not pre-adsorbed by ICV in this test. The results indicated a wide circulation in Africa of IDV, and its significant tropism to all the hosts studied.

H. Nedland et al. [23] examined 364 serum samples from horses of 141 farms in the Midwestern United States and showed that animals are susceptible to two IDV lines represented by D/swine/Oklahoma/1334/2011 (D/OK line) and D/bovine/Oklahoma/660/2013 (line D/660), as 12 % and 11 % of sera, respectively, were seropositive in HI.



The hosts of the influenza D virus. IDV can be transmitted between cattle and also between pigs as indicated by the solid line. Antibody specific for IDV were also detected in sera from small ruminants (sheep and goat), horses, camels and humans, especially in people with cattle exposure, but no virus was isolated (indicated by the broken line), indicating that IDV may be transmitted from cattle to humans (broken line)

M. Quast et al. [43] in a serological analysis of 648 sheep and goat blood serum samples collected in 2014 in the USA and Canada, detected antibodies to IDV in 5.2 % (29/557) of the studied sheep from 13.5 % (17/126) of farms. In turn, 8.8 % (8/91) of goats from 13.3 % (2/15) of the tested farms also contained antibodies against IDV.

IDV was first isolated from pigs and then from cattle. Since cattle seropositivity is much higher than that of pigs, it is believed that it is the main natural reservoir. Currently known IDV hosts are shown in figure. In addition, the ability of D/OK to infect ferrets, guinea pigs, and transmit them to native animals by direct contact is shown [26,44], which allows them to be used as models in the study of IDV.

Phylogenesis of influenza D. viruses. With the accumulation of IDV strains of various origins and places of isolation, the possibility of their comparative antigenic and phylogenetic studies appeared. E.A. Collin et al. [38] performed full genome sequencing and phylogenetic analysis of 7 viral RNA segments of six new strains and four previously registered IDVs and revealed two different circulating lines – D/OK and D/660, which often reassorted with each other. Antigenic analysis using representative viruses D/swine/Oklahoma/1334/2011 and D/bovine/Oklahoma/660/2013 and their antisera in the HI showed an approximately 10-fold loss of cross-reactivity. An important finding was that one of the isolates (D/bovine/Texas/3-13/2011) belonged to the D/bovine/Oklahoma/660/2013 cluster, but was characterized by high titers with a serum to the heterologous variant D/swine/Oklahoma/1334/2011. Molecular modeling of the HEF fusion protein of strain D/bovine/Texas/3-13/2011 made it possible to identify the mutation at position 212 responsible for unusual serological test titers. The obtained data indicate the widespread prevalence of at least two genetically different IDV lines in cattle, and also the vital role of lysine (K212) in antigenic recognition of D/swine/Oklahoma/1334/2011-like viruses. IDVs related to D/bovine/Oklahoma/660/2013 clade carried arginine in this position (R212).

Phylogenetic analysis of the HEF sequence of the USA IDV isolates performed by L. Ferguson et al. [26], aligned them into two genetic clusters: viruses D/bovine/Mississippi/C00046N/2014 and D/bovine/Mississippi/C00030P/2014 were genetically close to D/swine/Oklahoma/1334/2011 (D/OK), while D/bovine/Mississippi/C00013N/2014 and D/bovine/Mississippi/C00014N/2014 were genetically related to D/bovine/Oklahoma/660/2013 (D / 660).

O. Flynn et al. [39] found that five IDV strains isolated in Ireland in 2014–2016 were grouped in the same clade together with the virus from Europe D/swine/OK/1334/2011.

S. Murakami et al. [37] using reverse transcription-PCR successfully amplified the complete genome sequence of the first Japanese IDV strain (D/bovine/Ibaraki/7768/2016), sequenced its seven segments and built their phylogenetic trees. The results indicated that it occupies a separate position from strains from other countries, and only the M gene is included in one cluster with isolates from France (D/bovine/France/2986/2012) and China (D/bovine/Shandong/Y127/2014, D/bovine/Shandong/Y217/2014, D/bovine/Shandong/Y125/2014). Later on, in Japan, from a cow with signs of respiratory disease, the D/bovine/Yamagata/10710/2016 strain highly related D/bovine/Ibaraki/7768/2016 according to the HEF gene was isolated, the homology of the nucleotide and amino acid sequences was 99.8 % and 100 %, respectively [45]. H. Mekata et al. [41] performed analysis of 46 nasal swabs from cattle with signs of respiratory disease from 26 different farms in Japan, using the Next Generation Sequencing, the nucleotide sequence of the complete IDV genome was determined, and it was shown that it forms a separate cluster. According to the authors, the virus could develop uniquely over a long period, and its pathogenic properties differ from the strains found in other countries. In 2018 T. Odagiri et al. [46] studied IDV phylogenesis and established the presence of three genetically distinct lines: D/OK, D/660, and D/Japan. In addition, in the cross-linked HI, it was found that representatives of three lines in the composition of the surface protein HEF present both common for all and line-specific antigens.

The most conserved PB1 viral protein gene is often used to evaluate the evolutionary relationships of influenza viruses. S. Su et al. [2] based on the nucleotide sequences of the PB1 genes of influenza A, B, C, and D viruses, constructed a phylogenetic tree to determine the relationship between them. IDV clusters turned out to be the most closely related to ICV, which suggests their common ancestor. Sequence analysis of the PB2, P3, NP, M, and NS genes also confirmed the origin of IDV from human ICV [26]. To study the origin and evolutionary history of IDV, the authors conducted a Bayesian analysis of the HEF ICV gene sequences, indicating 1896 as the average time (t-MRCA) of the most recent common ancestor, which is consistent with previously obtained data [47]. For the HEF genes of ICV and

IDV, the t-MRCA value was 482 A.D. It is shown that two IDV lines – D/OK and D/660 had the last common ancestor (t-MRCA) about 44.6 years ago. The average frequency of substitution for the HEF IDV gene was calculated using Bayesian analysis and amounted to 1.54×10^{-3} , which exceeds the frequency of ICV [47]. From this point of view, the causative agent of influenza D has great epidemic potential, and in the future, it is necessary to monitor its development constantly.

Conclusion. To date, an intensive study of IDV has shown that it infects pigs, cattle, and small ruminants, but the full spectrum of its susceptible hosts remains to be determined. To better understand the causative agent of influenza D from epidemiological, pathological and biological characteristics, in particular, the ability to cause disease in humans and be transmitted from person to person, new, more in-depth studies using molecular genetic tools are required. A significant stock of broad species range of susceptible farm animals needs a study of IDV prevalence in Kazakhstan.

Taken together, the epidemiological and metagenomic data, as well as experimental infection of animals, showed that the influenza virus D is the causative agent of respiratory disease of cattle, and therefore there is the task of appropriate vaccination. Moreover, the antigenic heterogeneity of HEF IDV explains the need to take into account which strains of which line circulate in a given region to prepare the most effective vaccine against this infection.

Қ.Х. Жұматов, А.И. Қыдырманов, М.Х. Саятов

Микробиология және вирусология ғылыми өндірістік орталығы, Алматы, Қазақстан

ТҰМАУ D ВИРУСЫ – ORTHOMYXOVIRIDAE ТҰҚЫМДАСЫНДАҒЫ ЖАҢА ТҮР ҚАЛЫПТАСТЫРАТЫН ПАТОГЕНДЕР

Аңдатпа. Тұмау қоздырғыштары Orthomyxoviridae тұқымдасына жатады және А, В, С, D тұмауы болып, сондай-ақ Quaranjavirus, Thogotovirus және Isavirus туыстықтарына бөлінеді. D тұмауы вирусы алғаш рет 2011 ж. АҚШ-та шошқаның кеңсірік шайындысынан бөлініп алынды, кейінірек Франция, Қытай, Италия, Ирландия, Жапония, бірқатар Африка елдерінде ірі қара малдың арасында кең тарағаны, сонымен қатар оның күзендер мен теңіз шошқаларына жұғатыны анықталған. D тұмауы вирусына қарсы антиденелер жылқылардың, қой мен ешкілердің және малмен байланыста болған адамдардың қан сарысуынан кездеседі. D тұмауы вирусының РНҚ геномы 9 ақуыз синтезіне жауап беретін жеті фрагменттен тұрады. Ең ұзын үш сегмент PB2, PB1 және P3 полимеразаларын кодтайды, төртіншісі – HEF гемагглютинин-эстераза тұтасу ақуызы, бесіншісі – NP нуклеопротеині. Алтыншы фрагмент мембрана полипептидтерінің DM1 және DM2 синтезіне қатысады, олардың бірі вирустық мембрананы ішінен қаптап жатады, екіншісі протон арналарының қызметін орындайды. Жетінші сегмент NS1 құрылымдық емес ақуызды және NEP ядролық экспорт ақуызын кодтайды; NS1 жасушалық интерферонды бейтараптандыруға көмектеседі, NEP рибонуклеопротеиннің ядролық экспортын жүзеге асырады. Тұмау D вирусының үш – D/OK, D/660 және D/Жарап филогенетикалық желісі сипатталған, олар вакцина дайындау кезінде ескерілуі керек. Оның эпидемиологиялық, патологиялық және биологиялық сипаттамалары тұрғысынан, адамдарда ауру тудыруы және адамнан адамға берілуі мүмкін болатын қазіргі заманғы экологиялық-вирусологиялық және молекулалық-генетикалық әдістерді қолдана отырып, жаңа, тереңірек зерттеулер қажет деген тұжырым жасалды. Аса консервативті вирустық PB1 ақуыз гені тұмау вирустарының эволюциялық қатынастарын бағалау үшін жиі қолданылады. S. Su et al. [2] А, В, С және D вирустарының PB1 генінің нуклеотидтік тізбегі негізінде олардың арасындағы байланысты анықтау үшін филогенетикалық дарак құрастырылды. IDV кластерлері ICV-мен ең жақын байланыста болды, бұл олардың ата-бабаларының ортақ екенін білдіреді. PB2, P3, NP, M, NS гендерінің тізбегін талдау нәтижесінде, олардың адам ICV-нен IDV шыққанын растады [26]. IDV-ның пайда болуы мен эволюциялық тарихын зерттеу үшін авторлар HEF ICV гендік тізбегіне байесовский талдауын жүргізді, бұл 1896 ж., ортақ туыстастықтың орташа уақыты (t-MRCA) деп көрсетеді, бұл алдыңғы деректермен сәйкес келеді [47]. HEF гендері, ICV және IDV үшін t-MRCA мәні 482 н.т., болды. Екі линия IDV-D/OK және D/660 шамамен 44.6 жыл бұрын соңғы жалпы ата-бабалары (t-MRCA) болғанын көрсетеді. HEF IDV гені алмасуының орташа жиілігі Байес анализінің көмегімен есептелді және $1,54 \times 10^{-3}$ құрады, бұл ICV жиілігінен асып түсті [47]. Осы тұрғыдан алғанда, D тұмауының қоздырғышы үлкен эпидемиялық потенциалға ие және заманауи экологиялық-вирусологиялық және молекулалық-генетикалық әдістерді қолдану арқылы болашақта оның дамуын үнемі бақылау қажет.

Түйін сөздер: вирус, тұмау D, геном, өзгергіштік, мүйізді ірі қара, филогенез, клайд, тұтасу ақуызы HEF, серология.

К. Х. Жуматов, А. И. Кыдырманов, М. Х. Саятов

Научно производственный центр микробиологии и вирусологии, Алматы, Казахстан

ВИРУСЫ ГРИППА D – ПАТОГЕНЫ, ОБРАЗУЮЩИЕ НОВЫЙ РОД В СЕМЕЙСТВЕ ORTHOMYXOVIRIDAE

Аннотация. Возбудители гриппа относятся к семейству Orthomyxoviridae и разделяются на роды: Influenzavirus A, B, C, D, а также Quaranjavirus, Thogotovirus и Isavirus. Впервые вирус гриппа D выделили из назальных смывов свиней в 2011 г. в США, в дальнейшем показана его широкая распространенность среди крупного рогатого скота во Франции, Китае, Италии, Ирландии, Японии, ряде африканских стран, а также способность инфицировать хорьков, морских свинок. Антитела к вирусу гриппа D обнаружены в сыворотках крови лошадей, овец, коз и у людей, контактировавших с крупным рогатым скотом. РНК-геном вируса гриппа D представлен семью фрагментами, ответственными за синтез 9 белков. Самые длинные три сегмента кодируют полимеразы PB2, PB1 и P3, четвертый – белок слияния гемагглютинин-эстеразу HEF, пятый – нуклеопротеин NP. Шестой фрагмент участвует в синтезе мембранных полипептидов DM1 и DM2, один из которых выстилает вирусную мембрану изнутри, другой осуществляет функцию протонных каналов. Седьмой сегмент кодирует неструктурный белок NS1 и белок ядерного экспорта NEP; NS1 способствует нейтрализации клеточного интерферона, NEP опосредует ядерный экспорт рибонуклеопротеина. Описаны три филогенетические линии вируса гриппа D – D/OK, D/660 и D/Japan, что необходимо учитывать при приготовлении вакцин. Делается вывод о том, что с точки зрения его эпидемиологических, патологических и биологических характеристик, потенциальной способности вызывать заболевание у людей и передаваться от человека человеку требуются новые, более углубленные исследования. Наиболее консервативный ген вирусного белка PB1 часто используется для оценки эволюционных взаимоотношений вирусов гриппа. S. Su et al. [2] на основе нуклеотидных последовательностей генов PB1 вирусов гриппа A, B, C и D построили филогенетическое древо для определения взаимосвязи между ними. Кластеры IDV оказались наиболее тесно связаны с ICV, что предполагает их общего предка. Анализ последовательности генов PB2, P3, NP, M и NS также подтвердил происхождение IDV из ICV человека [26]. Чтобы изучить происхождение и эволюционную историю IDV, авторы провели байесовский анализ последовательностей генов HEF ICV, указав 1896 год как среднее время (t-MRCA) самого последнего общего предка, что согласуется с ранее полученными данными [47]. Для генов HEF ICV и IDV значение t-MRCA было 482 н.э. Показано, что две линии IDV – D/OK и D/660 имели последнего общего предка (t-MRCA) около 44,6 лет назад. Средняя частота замещения гена HEF IDV была рассчитана с использованием байесовского анализа и составила $1,54 \times 10^{-3}$, что превышает частоту ICV [47]. С этой точки зрения возбудитель гриппа D обладает огромным эпидемическим потенциалом, и в будущем необходимо постоянно следить за его развитием. использованием современных эколого-вирусологических и молекулярно-генетических методов.

Ключевые слова: вирус, грипп D, геном, изменчивость, крупный рогатый скот, филогенез, кластер, белок слияния HEF, серология.

Information about authors:

Zhumatov Kainar Kh., Doctor of Biol. Sci., Prof. LLP «SPC Microbiology and Virology», Almaty, Kazakhstan; kainar60@yahoo.com; <https://orcid.org/0000-0001-6312-5730>

Kydyrmanov Aidyn I., Doctor of Vet. Sci., LLP «SPC Microbiology and Virology», Almaty, Kazakhstan; aidyn.kydyrmanov@gmail.com; <https://orcid.org/0000-0002-8374-6128>

Sayatov Marat Kh., Doctor of Biol. Sci., Prof., Academician of NAS of the RK, LLP «SPC Microbiology and Virology», Almaty, Kazakhstan; sayatov37@mail.ru; <https://orcid.org/0000-0003-4740-9156>

REFERENCES

- [1] Mostafa A., Abdelwhab E.M., Mettenleiter T.C., Pleschka S. (2018) Zoonotic Potential of Influenza A Viruses A Comprehensive Overview, *Viruses*. 10(9): 497. DOI: 10.3390/v10090497 (in Eng.).
- [2] Su S., Fub X., Lia G., Kerlinc F., Veit M. (2017) Novel Influenza D virus: Epidemiology, pathology, evolution and biological characteristics. *Virulence*, Vol. 8, N 8, P. 1580–1591. <https://doi.org/10.1080/21505594.2017.1365216> (in Eng.).
- [3] Ejiri H., Lim C.K., Isawa H., Fujita R., Murota K., Sato T., Kobayashi D., Kan M., Hattori M., Kimura T., Yamaguchi Y., Takayama-Ito M., Horiya M., Posadas-Herrera G., Minami S., Kuwata R., Shimoda H., Maeda K., Katayama Y., Mizutani T., Saijo M., Kaku K., Shinomiya H., Sawabe K (2018) Characterization of a novel thogotovirus isolated from Amblyomma testudinarium ticks in Ehime, Japan: A significant phylogenetic relationship to Bourbon virus, *Virus Res.* Apr 2; 249: 57-65. DOI: 10.1016/j.virusres.2018.03.004 (in Eng.).

- [4] Cárdenas C., Ojeda N., Labra A., Marshall S.H. (2017) An updated proposal for classification of infectious salmon anemia virus strains, *Arch Virol. Sep*;162(9): 2861-2867. DOI: 10.1007/s00705-017-3440-z (in Eng.).
- [5] Allison A.B., Ballard J.R., Tesh R.B., Brown J.D., Ruder M.G., Keel M.K., Munk B.A., Mickley R.M., Gibbs S.E., Travassos da Rosa A.P., Ellis J.C., Ip H.S., Shearn-Bochsler V., Rogers M.B., Ghedin E., Holmes E.C., Parrish C.R., Dwyer C. (2015) Cyclic avian mass mortality in the Northeastern United States is associated with a novel orthomyxovirus, *Journal of Virology*, 89: 1389–403. DOI: 10.1128/JVI.02019-14 (in Eng.).
- [6] Webster R., Govorkova E. (2014) Continuing challenges in influenza *Ann. N.Y. Acad. Sci.*, 1323:115–139. DOI: 10.1111/nyas.12462 (in Eng.).
- [7] Sayatov M.H., Zhumatov K.H. (2005) Virusy grippa ptic i gripp A (H5N1) u cheloveka [Izv.NAN RK. Ser. biol. i med.] 2:3-9 (in Russ.).
- [8] Sayatov M.Kh., Kydyrmanov A.I., Zhumatov K.Kh. (2009) Antigennyi spektr virusov grippa, tsirkuliruiushchikh sredi dikikh i domashnikh ptits v Respublike Kazakhsta [Doklady NAN RK] 4:46-50 (in Russ.).
- [9] Zhumatov K.Kh. (2009) Gripp i ugroza vozniknoveniia novoi pandemii [Vestnik NAN RK] 3:38-42 (in Russ.).
- [10] Zhumatov K.Kh. (2014) Virusy grippa A: klassifikatsiia, struktura i rasprostranenie v biosphere [Izvestiia NAN RK. 3: 47-57 (in Russ.).
- [11] Kydyrmanov A.I., Sayatov M.Kh., Karamendin K.O., Zhumatov K.Kh., Asanova S.E., Daulbayeva K.D., Starick E., Fereidouni S. (2017) Monitoring of influenza A viruses in wild bird populations in Kazakhstan in 2002-2009 [Arch Virol] 162(1): 147-155. DOI: 10.1007/s00705-016-3076-4 (in Russ.).
- [12] Sayatov M.Kh., Zhumatov K.Kh., Kydyrmanov A.I., Karamendin K.O., Daulbaeva K.D., Asanova S.E., Kasymbekov Ye.T., Khan E.Ya., Suleimenova S.A. (2017) Monitoring virusov grippa A v dikoi ornitofaune Kazakhstana (2002-2015 gg.) [Doklady NAN RK] 2:130-136 (in Russ.).
- [13] Zhumatov K.Kh., Kydyrmanov A.I., Karamendin K.O., Daulbaeva K.D., Khan Ye.Ya., Kassymbekov Ye.T., Sayatov M.Kh., Fereidouni S. (2018) Phylogenetic analysis of genes of surface proteins of highly pathogenic influenza a subtype H5N1 virus, isolated from gull in Kazakhstan [Bulletin Of National Academy of Sciences of The Republic of Kazakhstan] 5:6–11. <https://DOI.org/10.32014/2018.2518-1467.1> (in Russ.).
- [14] Simonson L. (2004) Pandemic influenza and mortality: past evidence and projections for the future. Board on Global Health. The threat of pandemic influenza: are we ready? The National Academies Press; <http://www.nap.edu/catalog/11150.html> (in Eng.).
- [15] Asai N., Yokoi T., Nishiyama N., Koizumi Y., Sakanashi D., Kato H., et al. (2017) Secondary organizing pneumonia following viral pneumonia caused by severe influenza B: a case report and literature reviews, *BMC Infect Dis.*; 17:572. DOI: 10.1186/s12879-017-2677-1 (in Eng.).
- [16] Rota P.A., Shaw M.W., Kendal A.P. (1989) Cross-protection against microvariants of influenza virus type B by vaccinia viruses expressing haemagglutinins from egg- or MDCK cell-derived subpopulations of influenza virus type B/England/222/82, *J Gen Virol* 70(Part 6): 1533–1537. DOI:10.1099/0022-1317-70-6-1533 (in Eng.).
- [17] Chang C.P., New A.E., Taylor J.F., Chiang H.S. (1976) Influenza virus isolations from dogs during a human epidemic in Taiwan, *Int J Zoonoses* 3: 61–64. PMID: 977232 (in Eng.).
- [18] Kawano J., Onta T., Kida H., Yanagawa R. (1978) Distribution of antibodies in animals against influenza B and C viruses, *Jpn J Vet Res* 26: 74–80. PMID: 739713 (in Eng.).
- [19] Romvary J., Meszaros J., Barb K. (1980) Susceptibility of birds to type-B influenza virus, *Acta Microbiol Acad Sci Hung* 27: 279–287. PMID: 6258401 (in Eng.).
- [20] Osterhaus A.D., Rimmelzwaan G.F., Martina B.E., Bestebroer T.M., Fouchier R.A. (2000) Influenza B virus in seals, *Science*, 288(5468): 1051±3. PMID: 10807575 (in Eng.).
- [21] Zhiguang Ran, Huigang Shen, Yuekun Lang, Elizabeth A. Kolb, Nuri Turan, Laihua Zhu, Jingjiao Ma, Bhupinder Bawa, Qinfang Liu, Haixia Liu, Megan Quast, Gabriel Sexton, Florian Krammer, Ben M. Hause, Jane Christopher-Hennings, Eric A. Nelson, Juergen Richt, Feng Li, and Wenjun Ma. (2015). Domestic Pigs Are Susceptible to Infection with Influenza B Viruses, *J Virol*, 89(9): 4818–4826. DOI: 10.1128/JVI.00059-15 (in Eng.).
- [22] Guo Y.J., Jin F.G., Wang P., Wang M., Zhu J.M. (1983) Isolation of influenza C virus from pigs and experimental infection of pigs with influenza C virus, *J Gen Virol* 64: 177–182. <http://dx.doi.org/10.1099/0022-1317-64-1-177> (in Eng.).
- [23] Nedland H., Wollman J., Sreenivasan C., Quast M., Singrey A., Fawcett L., Christopher-Hennings J., Nelson E., Kaushik R.S., Wang D., Li F. (2018) Serological evidence for the co-circulation of two lineages of influenza D viruses in equine populations of the Midwest United States, *Zoonoses Public Health*. February; 65(1): e148–e154. DOI:10.1111/zph.12423 (in Eng.).
- [24] Daszak P., Cunningham A.A. & Hyatt A.D. (2000) Emerging infectious diseases of wildlife—threats to biodiversity and human health, *Science* 287: 443– 449. PMID: 10642539(in Eng.).
- [25] Jones K.E., Patel N.G., Levy M.A., Storeygard A., Balk D., Gittleman J.L., Daszak P. (2008) Global trends in emerging infectious diseases, *Nature* 451: 990– 993. DOI: 10.1038/nature06536 (in Eng.).
- [26] Hause B.M., Ducatez M., Collin E.A., Ran Z., Liu R., Sheng Z., Armien A., Kaplan B., Chakravarty S., Hoppe A.D., Webby R.J., Simonson R.R., Li F. (2013) Isolation of a novel swine influenza virus from Oklahoma in 2011 which is distantly related to human influenza C viruses, *PLoS Pathog.* 9:e1003176. <http://dx.doi.org/10.1371/journal.ppat.1003176> (in Eng.).
- [27] Hause B.M., Collin E.A., Liu R., Huang B., Sheng Z., Lu,W., Wang D., Nelson E.A., Li F. (2014) Characterization of a novel influenza virus in cattle and Swine: Proposal for a new genus in the orthomyxoviridae family, *mBio*. 5(2): e00031-14. DOI: 10.1128/mBio.00031-14 (in Eng.).
- [28] Andrew M.Q. King, Elliot J. Lefkowitz, Arcady R. Mushegian, Michael J. Adams, Bas E. Dutilh, Alexander E. Gorbalenya, Balázs Harrach, Robert L. Harrison, Sandra Junglen, Nick J. Knowles, Andrew M. Kropinski, Mart Krupovic, Jens H. Kuhn, Max L. Nibert, Luisa Rubino, Sead Sabanadzovic, Hélène Sanfaçon, Stuart G. Siddell, Peter Simmonds, Arvind Varsani, Francisco Murilo Zerbin, Andrew J. Davison (2018) Changes to taxonomy and the International Code of Virus

Classification and Nomenclature ratified by the International Committee on Taxonomy of Viruses, Archives of Virology 163: 2601–2631 <https://doi.org/10.1007/s00705-018-3847-1> (in Eng.).

[29] Ferguson L., Olivier A.K., Genova S., Epperson W.B., Smith D.R., Schneider L., Barton K., McCuan K., Webby R.J., Wan X.-F. (2016) Pathogenesis of influenza D virus in cattle, *J Virol* 90: 5636–5642. DOI:10.1128/JVI.03122-15 (in Eng.).

[30] Paterson D., Fodor E. (2012) Emerging Roles for the Influenza A Virus Nuclear Export Protein (NEP), *PLoS Pathog.*, 8, e1003019 (in Eng.).

[31] Song H., Qi J., Khedri Z., Diaz S., Yu H., Chen X., Varki A., Shi Y., Gao G.F. (2016) An Open Receptor-Binding Cavity of Hemagglutinin-Esterase-Fusion Glycoprotein from Newly-Identified Influenza D Virus: Basis for Its Broad Cell Tropism, *PLoS Pathog* 12(1): e1005411. DOI:10.1371/journal.ppat.1005411 (in Eng.).

[32] Langereis M.A., Zeng Q., Gerwig G.J., Frey B., von Itzstein M., Kamerling J.P., et al (2009) Structural basis for ligand and substrate recognition by torovirus hemagglutinin esterases, *Proceedings of the National Academy of Sciences of the United States of America*, 106 (37): 15897–902. DOI: 10.1073/pnas.0904266106 PMID: 19721004; PubMed Central PMCID: PMC2747215. (in Eng.).

[33] Ferguson L., Eckard L., Epperson W. B., Long L.P., Smith D., Huston C., Genova S., Webby R., Wan X.-F. (2015 December) Influenza D Virus Infection in Mississippi Beef Cattle, *Virology*. 486: 28–34. DOI:10.1016/j.virol.2015.08.030 (in Eng.).

[34] Jiang W.M., Wang S.C., Peng C., Yu J.M., Zhuang Q.Y., Hou G.Y., Liu S., Li J.P., Chen J.M. (2014) Identification of a potential novel type of influenza virus in Bovine in China, *Virus Genes*; 49: 493–6. <http://dx.doi.org/10.1007/s11262-014-1107-3> (in Eng.).

[35] Ducatez M.F., Pelletier C., Meyer G. (2011–2014) Influenza D virus in cattle, France, *Emerg Infect Dis*. 2015; 21: 368–71 (in Eng.).

[36] Chiapponi C., Faccini S., De Mattia A., Baioni L., Barbieri I., Rosignoli C., Nigrelli A., Foni E. (2016) Detection of influenza D virus among swine and cattle, Italy, *Emerg. Infect. Dis*. 22, 352–354. doi: 10.3201/eid2202.151439 (in Eng.).

[37] Murakami S., Endoh M., Kobayashi T., Takenaka-Uema A., Chambers J. K., Uchida K., Nishihara M., Hause B., Horimoto T. (August 2016) Influenza D Virus Infection in Herd of Cattle, Japan, *Emerging Infectious Diseases*. www.cdc.gov/eid. Vol. 22, N 8, 1517 (in Eng.).

[38] Collin E.A., Sheng Z., Lang Y., Ma W., Hause B.M., Li F. (2015) Cocirculation of two distinct genetic and antigenic lineages of proposed influenza D virus in cattle, *J Virol* 89:1036–1042. doi:10.1128/JVI.02718-14 (in Eng.).

[39] Flynn O., Gallagher C., Mooney J., Irvine C., Ducatez M., Hause B., McGrath G. and Ryan E. (2018) Influenza D virus in cattle, Ireland, *Emerg Infect. Dis*. 24: 389–391 (in Eng.).

[40] Horimoto T., Hiono T., Mekata H., Odagiri T., Lei Z., Kobayashi T., Norimine J., Inoshima Y., Hikono H., Murakami K., Sato R., Murakami H., Sakaguchi M., Ishii K., Ando T., Otomaru K., Ozawa M., Sakoda Y. and Murakami S. (2016) Nationwide distribution of bovine influenza D virus infection in Japan, *PLoS One* 11: e0163828 (in Eng.).

[41] Mekata H., Yamamoto M., Hamabe S., Tanaka H., Omatsu T., Mizutani T., Hause B.M., Okabayashi T. (2017 Nov 16) Molecular epidemiological survey and phylogenetic analysis of bovine influenza D virus in Japan, *Transbound Emerg Dis*. 2018 Apr;65(2):e355-e360. DOI: 10.1111/tbed.12765. Epub (in Eng.).

[42] Salem E., Cook E.A.J., Lbacha H.A., Oliva J., Awoume F., Aplogan G.L., Hymann E.C., Muloi D., Deem S.L., Alali S., Zouagui Z., Fèvre E.M., Meyer G., Ducatez M.F. (2017) Serologic evidence for influenza C and D virus among ruminants and camelids, Africa, 1991–2015. *Emerg. Infect. Dis*. 23: 1556–1559 (in Eng.).

[43] Quast M., Sreenivasan C., Sexton G., Nedland H., Singrey A., Fawcett L., Miller G., Lauer D., Voss S., Pollock S., Cunha C.W., Christopher-Hennings J., Nelson E., Li F. (2015) Serological evidence for the presence of influenza D virus in small ruminants, *Vet. Microbiol*. 180: 281–285 (in Eng.).

[44] Sreenivasan C., Thomas M., Sheng Z., Hause B.M., Collin E.A., Knudsen D.E., Angela Pillatzki, Eric Nelson, Dan Wang, Radhey S. Kaushik, Feng Li. (2015) Replication and transmission of novel bovine influenza D virus in guinea pig model, *Journal of virology*. DOI: 10.1128/JVI.01630-15 PMID: 26378161 (in Eng.).

[45] Nakatsu S., Murakami S., Shindo K., Horimoto T., Sagara H., Noda T. and Kawaoka Y. (2018) Influenza C and D viruses package eight organized ribonucleoprotein complexes, *J. Virol*. 92: e02084–e17 (in Eng.).

[46] Odagiri T., Ishida H., Li Jun-You, Endo M., Kobayashi T., Kamiki H., Matsugo H., Takenaka-Uema A., Murakami S., T. (2018) Horimoto. Antigenic heterogeneity among phylogenetic clusters of influenza D viruses, *J. Vet. Med. Sci*. 80 (8): 1241–1244, DOI: 10.1292/jvms.18-0157 (in Eng.).

[47] Gatherer D. (2010) Tempo and mode in the molecular evolution of influenza C, *PLoS Currents*; 2:RRN1199. PMID:21127722 (in Eng.).

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www.nauka-nanrk.kz

ISSN 2518-1629 (Online), ISSN 2224-5308 (Print)

<http://biological-medical.kz/index.php/en/>

Редакторы: *М. С. Ахметова, Г. Б. Халидуллаева, Д. С. Аленов*
Верстка на компьютере *Д. А. Абдрахимовой*

Подписано в печать 02.04.2020.

Формат 60x881/8. Бумага офсетная. Печать – ризограф.
5,5 п.л. Тираж 300. Заказ 2.